

## DOCTOR OF CLINICAL PSYCHOLOGY (DCLINPSY)

### Doctorate in Clinical Psychology: Main Research Portfolio

**1) Critical Literature Review: A model of psychological adjustment to scarring following planned surgery; 2) Service Improvement Project: Volunteers' experiences of helping hoarders and hoarders' experiences of being helped; 3) Main Research Project: Post-traumatic growth in mothers 2 - 6 years after stillbirth and early miscarriage.**

Ryninks, Kirsty

*Award date:*  
2018

*Awarding institution:*  
University of Bath

[Link to publication](#)

### Alternative formats

If you require this document in an alternative format, please contact:  
[openaccess@bath.ac.uk](mailto:openaccess@bath.ac.uk)

Copyright of this thesis rests with the author. Access is subject to the above licence, if given. If no licence is specified above, original content in this thesis is licensed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International (CC BY-NC-ND 4.0) Licence (<https://creativecommons.org/licenses/by-nc-nd/4.0/>). Any third-party copyright material present remains the property of its respective owner(s) and is licensed under its existing terms.

#### Take down policy

If you consider content within Bath's Research Portal to be in breach of UK law, please contact: [openaccess@bath.ac.uk](mailto:openaccess@bath.ac.uk) with the details. Your claim will be investigated and, where appropriate, the item will be removed from public view as soon as possible.

# Research Portfolio Submitted in Part Fulfilment of the requirements for the Degree of Doctorate in Clinical Psychology

---

Kirsty Emma Ryninks

Doctorate in Clinical Psychology

University of Bath  
Department of Psychology

May 2018

## **COPYRIGHT**

Attention is drawn to the fact that copyright of this thesis rests with the author. A copy of this thesis has been supplied on condition that anyone who consults it is understood to recognise that its copyright rests with the author and that they must not copy it or use material from it except as permitted by law or with the consent of the author.

## **RESTRICTIONS ON USE**

This thesis may be made available for consultation within the University Library and may be photocopied or lent to other libraries for the purposes of consultation with effect from .....

Signed on behalf of the Faculty / School of .....



## **Word Counts**

*Note, all word counts exclude abstracts, figures and tables*

<b>Critical Literature Review</b>	7017
-----------------------------------	------

A Model of Psychological Adjustment to Scarring following Planned Surgery

<b>Service Improvement Project</b>	5189
------------------------------------	------

Volunteers' experiences of helping hoarders and hoarders' experiences of being helped

<b>Main Research Project</b>	5318
------------------------------	------

Posttraumatic growth in mothers 2-6 years after stillbirth and early miscarriage

<b>Executive Summary</b>	958
--------------------------	-----

<b>Connecting Narrative</b>	2968
-----------------------------	------





## Table of Contents

Word Counts .....	3
Table of Tables.....	9
Table of Figures .....	10
<b>Abstracts</b> .....	11
Critical Literature Review.....	11
Service Improvement Project.....	12
Main Research Project .....	13
<b>A Model of Psychological Adjustment to Scarring following Planned Surgery</b> .....	15
Outline of the Review .....	17
Scarring .....	17
Understanding Disfigurement.....	19
Surgical Treatments and Scarring.....	20
Psychological Preparation for Scarring following Surgery .....	24
Models of Adjustment, Disfigurement and Appearance-Related Disorders.....	25
The ARC Framework of Adjustment to Visible Difference (Clarke, Thompson, Jenkinson, Rumsey, & Newell, 2014) .....	26
A Cognitive Behavioural Model of Body Dysmorphic Disorder (BDD) .....	27
A New Understanding of Psychological Adjustment to Scarring following Planned Surgery.....	29
A Model of Psychological Adjustment to Scarring following Planned Surgery .....	29
Before the Operation/Surgery.....	29
After the Operation/Surgery .....	32
Summary .....	34
Clinical Implications.....	34
References.....	36
<b>Volunteers' experiences of helping hoarders and hoarders' experiences of being helped</b> .....	45
Introduction.....	47
Method .....	49
Participants and Recruitment.....	49
Design.....	49
Data collection.....	50
Analysis .....	50
Results.....	51
Sample Characteristics .....	51

Qualitative Analysis .....	52
Discussion.....	58
Conclusions and Implications .....	61
References .....	62
<b>Posttraumatic growth in mothers 2-6 years after stillbirth and early miscarriage .....</b>	<b>65</b>
Introduction .....	67
Method.....	70
Design .....	70
Participants.....	70
Procedure.....	70
Materials.....	71
Statistical Analysis .....	73
Results .....	73
Participant Characteristics.....	73
Descriptive statistics.....	75
Variables predicting PTG.....	76
Discussion.....	79
Strengths and Limitations.....	80
Clinical Implications .....	82
Conclusions .....	82
References .....	83
<b>Executive Summary .....</b>	<b>91</b>
<b>Connecting Narrative .....</b>	<b>95</b>
Overview .....	95
Case studies .....	95
Service Improvement Project .....	97
Main Research Project and Critical Review of Literature .....	98
Critical Review of Literature.....	98
Main Research Project .....	99
Summary.....	101
<b>Acknowledgments .....</b>	<b>103</b>
<b>Appendices.....</b>	<b>105</b>
Appendix I: Instructions for Authors – Body Image .....	105
Appendix II: Copyright permission (The ARC Framework of Adjustment to Visible Difference) - Wiley.....	119

Appendix III: Copyright permission (The cognitive behavioural model of Body Dysmorphic Disorder) – David Veale.....	120
Appendix IV: Instructions for Authors – British Journal of Clinical Psychology .....	121
Appendix V: University of Bath Service Improvement Project Ethical Approval .....	125
Appendix VI: Service Improvement Project Study Materials (Client and volunteer/case worker: Information Sheet, Consent Form, Demographic Questionnaires, Debrief Sheet) .....	126
Appendix VII: Service Improvement Project Outcomes .....	138
Appendix VIII: Instructions for Authors –BMC Pregnancy and Childbirth.....	139
Appendix IX: Permission to use questionnaires from Posttraumatic Growth Research Centre .....	146
Appendix X: University of Bath Main Research Project Ethical Approval .....	147
Appendix XI: Main Research Project Study Materials (Information Sheet, Consent Form, Demographic Questionnaire, Study Questionnaires, Debrief Sheet) .....	148



## Table of Tables

### Service Improvement Project

Table 2.1. <i>Sample Characteristics (N = 11)</i> .....	51
---	----

Table 2.2. <i>Superordinate and Subordinate themes</i> .....	52
--	----

### Main Research Project

Table 3.1 <i>Sample Characteristics (N = 120)</i> .....	74
---	----

Table 3.2. <i>Means (and standard deviations) for questionnaire scores by type of loss</i> .....	75
--	----

Table 3.3. <i>Correlations for SB and EM groups</i> .....	77
---	----

Table 3.4. <i>Multiple Regression Analysis: variables included in the regression model</i> .....	78
--	----

## **Table of Figures**

### **Critical Literature Review**

<i>Figure 1.1. The ARC Framework of Adjustment to Visible Difference.....</i>	<i>26</i>
<i>Figure 1.2. The Cognitive Behavioural model of BDD.....</i>	<i>28</i>
<i>Figure 1.3. A Model of Psychological Adjustment to Scarring following Planned Surgery.....</i>	<i>30</i>

## **Abstracts**

### **Critical Literature Review**

Individuals' lives can be profoundly affected by an altered appearance (Moss, 2005) and the perception of altered physical appearance may disturb how individuals perceive and value their bodies. Evidence suggests that socially awkward and avoidant behaviour is often displayed by others in the context of individuals with a disfigured appearance (Bull & Rumsey, 2012). 4.7 million people were admitted to hospital for a surgical procedure in the UK in 2013/14 (Royal College of Surgeons, 2016). While most individuals are happy to have treatment to improve their medical conditions and/or lengthen their life, many also face the reality of altered appearance through scarring as a result of their surgery. There may be limited understanding in the medical field that what is done in surgery in the context of "fixing" could have the impact of creating another set of problems for patients. Scars rarely pose a health risk and yet patients frequently present with scar-related aesthetic, social and psychological distress (Chacon, França, Ledon, & Nouri, 2013). Although a number of models have been developed to understand the concepts of appearance and disfigurement, there is limited research which directly addresses the important 'side effect' of common medical procedures, 'surgical scarring'. This narrative review aims to synthesise the existing literature on surgical scarring and appearance within a broadly cognitive-behavioural framework to develop a better understanding of psychological adjustment to scarring following planned surgery. A number of existing models provide a useful framework to understand adjustment to scarring following burns, traumatic injury and pre-existing disfiguring conditions; however, they do not sufficiently aid understanding of adjustment to planned surgery and health professionals do not have a framework to aid targeted preparation for surgery/treatment. We therefore propose a new model of psychological adjustment to scarring following planned surgery and focus on aiding health professionals to better help their patients through the development of specific and targeted preparation and treatment in the future. The review concludes by highlighting the importance of shifting the focus in medicine from 'functioning body parts' to positive psychological adjustment and wellbeing following planned surgery.



## **Service Improvement Project**

**Objectives.** The present study aimed to explore the experiences of compulsive hoarders and volunteer helpers within the context of a UK-based charity providing support to older adults with hoarding difficulties.

**Design.** Qualitative methods were adopted to investigate the lived experience of participants and add meaning to current understandings of compulsive hoarding.

**Methods.** A total of 7 volunteers and 4 clients (compulsive hoarders) were recruited and interviewed using a semi-structured interview, designed to explore experiences of providing and receiving help. Qualitative analysis of the interview data was performed using Interpretive Phenomenological Analysis.

**Results.** Four superordinate themes were identified: relationship between client and volunteer; ‘live life again’; challenges; and supporting volunteers. The relationship was crucial in providing a trusting foundation from which clients felt able to move forward. Volunteers provided a space for clients to talk and appropriate self-disclosure helped to build a relationship. The informal and ‘non-professional’ status of volunteers enabled clients to take the lead and feel more in control of the therapeutic process. Volunteer flexibility and lack of time constraints contributed to clients ‘making space’. The support from volunteers enabled clients to ‘live life again’ and created a domino effect, bringing about improvements in other areas of their lives.

**Conclusions.** The findings will be discussed in relation to the training of health professionals to work with people with hoarding difficulties and the implications of the findings for treatment approaches and service provision.

## **Practitioner Points**

- Switching the focus from hoarding and associated behaviours to the client as a person and their interests may create a space for change and bring about improvements in compulsive hoarders’ quality of life.
- Services may benefit from developing separate pathways for compulsive hoarders, focused on allowing volunteers and professionals time and flexibility to build the therapeutic relationship in the future.
- In light of this study’s small sample size, the findings may have limited generalisability.
- Purposive sampling may have excluded clients experiencing greater shame and embarrassment.

## **Main Research Project**

**Background.** The possibility of posttraumatic growth (PTG) in the aftermath of pregnancy loss has received limited attention to date. This study investigated PTG in mothers two to six years following stillbirth (SB) compared to early miscarriage (EM). It was hypothesised that mothers following SB will demonstrate more (1) PTG, (2) challenge to assumptive beliefs, and (3) disclosure than mothers following EM. The study also sought to understand how theoretically-derived variables of the Model of Growth in Grief (challenge to assumptive beliefs and disclosure) explained unique variance in PTG when key factors were controlled for.

**Methods.** One hundred and twenty women who had experienced a SB (N=57) or EM (N=63) two to six years ago completed validated questionnaires relating to PTG and key variables relevant to emotional adjustment post-bereavement.

**Results.** Participants who had experienced a SB demonstrated significantly higher levels of PTG, posttraumatic stress symptoms, perinatal grief, disclosure, challenge to assumptive beliefs and rumination than participants who had experienced an EM. In a hierarchical stepwise regression analysis, challenge to assumptive beliefs alone predicted 17.6% of the variance in PTG. Intrusive and deliberate rumination predicted an additional 5.9% of variance, with urge to talk and actual self-disclosure predicting a further 14.7%. A final model including these variables explained 46.4% of the variance in PTG.

**Conclusions.** Significantly higher levels of PTG were found in mothers following SB compared to in mothers following EM. Mothers experienced greater challenge to their assumptive beliefs and revealed higher levels of actual self-disclosure and greater urge to talk following their SB. These findings can partially be explained by differences in key variables from the Model of Growth in Grief.

**Keywords:** Posttraumatic growth, stillbirth, early miscarriage, disclosure, pregnancy loss



**A Model of Psychological Adjustment to Scarring following Planned Surgery**

Kirsty Ryninks

Doctorate in Clinical Psychology

Department of Psychology, University of Bath, Claverton Down, Bath, BA2 7AY

Email: K.E.Ryninks@bath.ac.uk

Word count: 7017

May 2018

**Internal/academic supervisors:** Professor Paul Salkovskis and Dr Cara Davis;

Doctoral Programme in Clinical Psychology

Department of Psychology, University of Bath, Claverton Down, Bath, BA2 7AY

Email: P.M.Salkovskis@bath.ac.uk and C.Davis@bath.ac.uk

**External/field supervisor:** Dr Nell Ellison, Bristol Royal Hospital for Children

University Hospitals Bristol NHS Foundation Trust

Upper Maudlin Street, Bristol, BS2 8BJ

Email: nell.ellison@uhbristol.nhs.uk

**Proposed Journal:** Body Image

Please see Appendix I for instructions to authors



## **Outline of the Review**

Individuals' lives can be profoundly affected by an altered appearance (Moss, 2005) and the perception of altered physical appearance may disturb how individuals perceive and value their bodies. Evidence suggests that socially awkward and avoidant behaviour is often displayed by others in the context of individuals with a disfigured appearance (Bull & Rumsey, 2012). 4.7 million people were admitted to hospital for a surgical procedure in the UK in 2013/14 (Royal College of Surgeons, 2016). While most individuals are happy to have treatment to improve their medical conditions and/or lengthen their life, many also face the reality of altered appearance through scarring as a result of their surgery. There may be limited understanding in the medical field that what is done in surgery in the context of "fixing" could have the impact of creating another set of problems for patients. Scars rarely pose a health risk and yet patients frequently present with scar-related aesthetic, social and psychological distress (Chacon, França, Ledon, & Nouri, 2013). Although a number of models have been developed to understand the concepts of appearance and disfigurement, there is limited research which directly addresses the important 'side effect' of common medical procedures, 'surgical scarring'.

This narrative review aims to synthesise the existing literature on surgical scarring and appearance within a broadly cognitive-behavioural framework to develop a better understanding of psychological adjustment to scarring following planned surgery. A number of existing models provide a useful framework to understand adjustment to scarring following burns, traumatic injury and pre-existing disfiguring conditions; however, they do not sufficiently aid understanding of adjustment to planned surgery and health professionals do not have a framework to aid targeted preparation for surgery/treatment. We therefore propose a new model of psychological adjustment to scarring following planned surgery and focus on aiding health professionals to better help their patients through the development of specific and targeted preparation and treatment in the future. The review concludes by highlighting the importance of shifting the focus in medicine from 'functioning body parts' to positive psychological adjustment and wellbeing following planned surgery.

## **Scarring**

Scars are the body's natural response to injury or damage and are areas of fibrosis that occur in normal skin following injury (França, Chacon, Ledon, Savas, & Nouri, 2013). They are the imperfect end point of the normal continuum of tissue repair (Bayat, McGrouther, & Ferguson, 2003) and can be classified as normotrophic, hypertrophic and atrophic scars (Chacon et al., 2013). Whilst hypertrophic and keloid scars are both elevated

and raised, keloids grow beyond wound boundaries and are usually pink-red, firm, itchy and tender (Philipp, Scharschmidt, & Berlien, 2008). Both keloid and hypertrophic scars can result in pain but keloids cause more severe symptoms including physical discomfort and burning sensations (Lupton & Alster, 2002).

Although scarring is a natural part of the healing process, living with scars can be challenging in a social and cultural environment that places high value on physical attractiveness and unblemished appearance (Beuf, 1990). Meaning is key; scar patients can experience wide-ranging effects that can have a major influence on their function, cosmetic appearance and psychological wellbeing (Brown, McKenna, Siddhi, McGrouther, & Bayat, 2008; França et al., 2013). Whilst for many individuals the scar may represent a symbol of recovery, a badge of honour or battle scar that can be viewed with pride, for others, the scar may be a constant reminder of what they have been through and source of stigma (Williamson & Wallace, 2012). King, McFetridge-Durdle, LeBlanc, Anzarut, and Tsuyuki (2009) reported that women's cardiac scars served as a reminder that they were: 'very ill and possibly close to death; had been through a great ordeal (either the surgery itself or beyond the surgery); had a defect corrected or remain with a weakness; and needed to take care to avoid more surgery' (King et al., 2009, p. 115).

The majority of published literature on scarring has focused on burns and traumatic injuries and therefore the management of patients with burn scars has been well documented in the literature. Across all age groups, 70% of patients with burn injury are male (Rajipura, 2002) and evidence suggests that psychological and psychiatric disorders are overrepresented in individuals with burn injuries (Klinge, Chamberlain, Redden, & King, 2009). Van Loey and Van Son (2003) reported that burn scars can have long-lasting psychological and physical implications, even after seeking remedial or cosmetic treatment, and the trauma of the accident and subsequent painful treatment may in fact induce the psychopathological responses in the first place. Lawrence, Mason, Schomer, and Klein (2012) conducted a systematic review of the impact of scarring after burn injury, indicating that psychosocial variables (e.g. social comfort and perceived stigmatisation) were more highly correlated with body image than scar characteristics. Therefore, in addition to psychological vulnerabilities contributing to the occurrence of a burn injury, psychological factors are also involved in the harm and impact that a burn injury can cause. Moreover, burn severity is not always predictive of psychological adjustment (Tedstone & Tarrier, 1997) and small burns have been reported to cause strong psychological distress (Shakespeare, 1998).

## **Understanding Disfigurement**

Living with disfigurement can have a profound effect on those affected and intuitively, greater objective severity of a disfiguring condition may be expected to cause more psychosocial distress (Brown, Moss, McGrouther, & Bayat, 2010). In other words, an individual with extensive scarring may be expected to experience greater difficulty than someone with fewer scars (Moss, 2005). Although there is evidence of this position from anecdotal accounts (Bernstein, 1982; Thomas, 1990), MacGregor (1970) noted in her clinical work that some individuals with mild disfigurement had more difficulty adjusting than individuals with more severe disfigurements. In addition, individuals with no observable disfigurement (e.g. in Body Dysmorphic Disorder) often report high levels of distress. A number of studies in a variety of appearance conditions have since reported this position, with no clear relationship between objective severity or physical extent of the disfigurement and adjustment or level of psychosocial distress (Rumsey, Clarke, White, Wyn-Williams, & Garlick, 2004; Sarwer, Whitaker, & Bartlett, 2001; Thompson, Kent, & Smith, 2002).

Moss (2005) investigated subjective and objective severity of disfigurement and psychological adjustment in four hundred surgical patients with appearance altering conditions. Patients subjectively rated the severity of their disfigurement and objective ratings of severity were made by their plastic surgeons. Results revealed that poorer adjustment was related to greater subjectively perceived severity of disfigurement. However, assessment of perceived appearance severity was measured using a simple likert scale and the ratings of subjective and objective severity were completed on two separate patient groups (objective ratings by plastic surgeons for the outpatient group and subjective ratings by patients in the waiting list group). The authors of the study therefore concluded that future research should replicate the findings within the same sample and with more advanced measures of severity. Despite there being some limitations to the study, the findings provide empirical evidence to support previously anecdotal reports that objective ratings of disfigurement do not necessarily predict distress. Rumsey and Harcourt (2004) also described anecdotal observations of the linear relationship between adjustment and subjective severity of disfigurement in their review paper on body image and disfigurement.

Brown et al. (2010) later examined psychosocial distress in patients attending a specialist scar plastic surgery clinic. Skin scars were independently assessed using validated measures on two occasions, overcoming the limitations of previous research. The results demonstrated that patients with non-visible scars suffered more psychological distress than those with visible scars, as non-visible scars provide limited chances for exposure and



conditioning. In addition, patients' own subjective assessment of their scar severity influenced patients' psychosocial morbidity ( $r = 0.48, p < 0.0001$ ), whilst objective scar severity ( $r = 0.14, p = 0.11$ ) did not. Evidence therefore suggests that psychological distress is not related to clinician-rated scar severity or scar type but patient-perceived scar severity is related. Although some might regard this as counter-intuitive, it appears to be the patient's subjective opinion, not objective rating, that is important when considering adjustment. This fits well with a cognitive perspective.

### **Surgical Treatments and Scarring**

Chacon et al. (2013) reported that an estimated 100 million people in developed nations each year will acquire scars following a surgical procedure. Keloids and hypertrophic scars are the most common types of scarring following surgery and although they are not medically dangerous, they are often painful and may result in significant disfigurement to the affected area (Gauglitz, Korting, Pavicic, Ruzicka, & Jeschke, 2011). Most surgical treatments have the potential to alter appearance and result in skin scarring or disfigurement (Williamson & Wallace, 2012), including head and neck cancer (facial surgery), abdominal stoma, prostate cancer, amputation, transplants, breast cancer (lumpectomy and mastectomy) and a number of life-saving cardiac treatments. Psychological adjustment and reactions to scarring following these treatments are less well understood than in burn injury and therefore a brief summary of the literature on three of the most common conditions where surgical treatments are required will be presented next.

#### ***Head and Neck Cancer***

As the face is considered a psychologically important area of the body, given its visibility and the value society places on facial attractiveness, any disfigurement to the facial area can often lead to severe consequences in social situations (Tebble, Thomas, & Price, 2004; Versnel, Plomp, Passchier, Duivenvoorden, & Mathijssen, 2012). Treatment in head and neck cancer is focused on the throat, mouth, ear, nose and eye regions and results in a number of difficulties for patients, including scarring. Appearance change is expected and treatments for advanced stages of the disease dramatically alter appearance (Williamson & Wallace, 2012).

Fingeret et al. (2012) reported that 75% of patients who had surgery as part of their treatment reported feeling concerned or embarrassed by changes to their body. Whilst this percentage is significant, the authors cautioned the generalisability of their findings as participants received treatment at a comprehensive cancer centre where the cases of head and neck cancer may have been more unusual and aggressive. However, other studies

highlight the importance of subjective ratings when considering adjustment. Negative impact on self-image is also common (Gamba et al., 1992) and Piff (1998) reported that patients often grieve the loss of their previous appearance following treatment. Social isolation is a usual occurrence in head and neck cancer and some patients have reported feeling stigmatised, with difficulty making friends and securing employment following treatment (Kent, 2000).

Anxiety, depression and reduced wellbeing have been well documented in the head and neck cancer population following surgery and Semple, Sullivan, Dunwoody, and Kernohan (2004) in their review article stated that depression and anxiety were reported by 30% to 40% of patients. There is also evidence that gender is implicated in adjustment in head and neck cancer, with women reporting lower life happiness and greater depressive symptoms (Katz, Irish, Devins, Rodin, & Gullane, 2003) than men following surgery. Social support appeared to act as a buffer between greater disfigurement and wellbeing in women, not men (Katz, Irish, Devins, Rodin, & Gullane, 2003), suggesting that women with limited social support may be more at risk of poor adjustment following surgery. In addition, greater appearance dissatisfaction and concern post-surgery have been reported in women compared to men (Liu, 2008). It has been suggested that this may be attributed to the higher importance of facial attractiveness in women, as well as the head and neck area being the most individual and personal part of the body (Williamson & Wallace, 2012). However, as only 18% of the possible sample returned their questionnaires (Liu, 2008), the findings should be interpreted with caution. Other findings by Katz et al. (2003) suggest that not all patients struggle with their appearance following treatment. Moreover, healthcare professionals are often left surprised by the distress caused from minimal changes in appearance in light of what they considered a ‘successful aesthetic outcome’. Psychological distress and adjustment therefore cannot be predicted from the type or objective severity of scarring and disfigurement in head and neck cancer, as it is often the individual’s subjective assessment that determines psychological wellbeing.

### ***Breast Cancer***

Treatments in breast cancer have advanced in recent decades and an increasing number of women are living with an altered appearance as a result of lumpectomies (surgical removal of part of the breast) and mastectomies (surgical removal of the whole breast). At a time of adjustment to diagnosis, women are required to make prompt decisions about their treatment and evidence suggests that appearance and body image are important factors influencing women’s decisions (Williamson & Wallace, 2012). Alfano and Rowland (2006) in their review article reported that women can describe mastectomy as

a psychologically painful experience, threatening their body image and sense of identity, which is not surprising given the social connotations of femininity, sexuality and motherhood associated with women's breasts (Khan et al., 2000). In a qualitative study with nineteen women, patients frequently reported difficulties with intimacy following breast cancer surgery and attempted to conceal their scars for fear of stigma or shocking others (Piot-Ziegler, Sassi, Raffoul, & Delaloye, 2010).

Older women tend to report more positive body image following surgery than younger women, suggesting that age may also be involved in appearance-related distress, with younger women more likely to experience problems (Nissen et al., 2001). Figueiredo, Cullen, Hwang, Rowland and Mandelblatt (2004) recruited a large sample of 563 women and focused on understanding the experiences of older women (aged 67 or older), a group that had been overlooked in previous studies. Women were more likely to experience distress and poorer adjustment to their altered body if they highly valued physical appearance of their breasts (Figueiredo, Cullen, Hwang, Rowland, & Mandelblatt, 2004). Although participants were interviewed at three time points, increasing understanding of changes in adjustment over the time, the authors acknowledged that the findings may have been different if a pre-surgical assessment had been conducted (as all interviews were conducted post-surgery). Body image also proved an important factor in 31% of women's decisions about their breast cancer treatment (Figueiredo, Cullen, Hwang, Rowland, & Mandelblatt, 2004). Fobair et al. (2006) found that half of the 548 young women with breast cancer in their study experienced problems with their body image some or much of the time following treatment. Fifty five percent of women reported feeling embarrassed by their body and 45% reported feeling less feminine. Although the study only focused on young women aged 50 or under, the sample included 29% of women of colour, overcoming the limitations of earlier studies with predominantly white and well-educated samples. Feelings of emptiness and grief have also been reported by women when their changed body is compared to their former self (Alfano & Rowland, 2006). Therefore, as reported in head and neck cancer, it is the subjective satisfaction with appearance that appears to determine psychological outcome following surgical treatment for breast cancer (Williamson & Wallace, 2012).

### ***Cardiac Surgery***

Whilst the psychological impact and management of scars in head and neck and breast cancer has received some attention in the literature, very little research has been reported in psychological adjustment and reactions to scarring following life-saving cardiac surgery. The most visible sign of the surgical intervention in cardiac treatment is the scar

and therefore although one would expect body image to be given attention, Masi and Brovedani (1999) reported that few findings have in fact been published. This continues to be the case today. Body image and experience of surgical scars are important factors in psychological adjustment following cardiac surgery. In a study of 201 patients with a scar from congenital heart disease surgery, 22-25% of individuals experienced negative psychological effects of their surgical scars (Crossland et al., 2005). Geyer et al. (2006) conducted standardised interviews and medical examinations with 343 patients with congenital cardiac disease and found that the outward appearance of many patients, especially the scar and any physical impairments present, were reminders that their heart had been surgically corrected (Geyer et al., 2006). Geyer et al. (2006) therefore reported that an unfavourable body image may lead individuals to perceive themselves as defective, in turn resulting in self-doubts and a sense of insecurity. Although the study's authors cautioned their findings' generalisability to patients with more severe congenital heart disease, evidence from other studies highlight the importance of subjective ratings when considering adjustment.

King et al. (2009) explored the impact of sternal scar formation in thirteen women and found that women were not well-prepared to learn to live with their scar and were left feeling less attractive. They identified a solitary process of 'coming to terms' and 'learning to live' with their scar and some women worried that others would judge them for 'not taking care of their health', possibly due to the connotations of lifestyle choices and acquired cardiac problems. Women were also negatively affected by their subjective perception of the cosmetic impact of the scar and this effect extended months and years after their surgery. However, as the interviews were conducted at one time point (1.5-3.75 years post-surgery), it is not possible to determine how perceptions of scarring changed over time. Cardiac surgery scars were also associated with lower self-esteem and body image in one hundred adult patients with Congenital Heart Disease (Kantoch et al., 2006). Forty eight percent of patients reported that they try to conceal their scar in public and more than half of patients believed that their scars permanently disfigure their bodies. Perception of chest scarring was the same regardless of when scars were acquired, although scars acquired in adulthood exerted less impact on self-esteem and self-confidence. A greater proportion of women than men felt that their scars negatively impacted their self-confidence and self-esteem (Kantoch et al., 2006). The authors noted that the impact of past events and patients' congenital heart disease on questionnaire responses in relation to the impact of their scar is unknown. However, as reported previously, it is the patient's subjective experience of the scar itself that counts.

## **Psychological Preparation for Scarring following Surgery**

The management of scars presents a challenge for health professionals, as scarring can greatly impact quality of life, with proneness to shame, body image problems and lack of self-esteem (Bock, Schmid-Ott, Malewski, & Mrowietz, 2006; França et al., 2013). Psychological preparation for surgery may help to reduce this challenge and patient information and preparedness have long been considered key elements of enhanced recovery pathways (Grocott, Martin, & Mythen, 2012). Whilst there are British Psychological Society (BPS) guidelines around preparation for undergoing medical procedures in children (Gaskell, 2010), there is no official guidance regarding psychological preparation for the post-procedural consequences of interventions in children or adults.

There continues to be a lack of published research addressing the important question of psychological preparation for surgery and studies continue to report poor psychological outcomes in patients following surgical scarring. King et al. (2009) in their qualitative study reported that patients were not ready for, or did not understand what was going to happen to their body following cardiac surgery. Evidence of poor psychological outcomes from a number of other studies led Papaspyros, Patel, Polyzois, Javagula, and Jeffrey (2011) to conclude that the potential impact of scar formation should be discussed with patients during consultation for surgery in their review of scar formation and impact on patient quality of life.

Patient information has traditionally been delivered in surgical and anaesthetic pre-assessment clinics, although there is increasing evidence for more sophisticated psychological preparation for surgery. Hood (2010) developed a body image toolkit for patients who experience a change to their appearance through their illness and cancer treatment. The toolkit was part of a project to improve supportive care and included a cognitive-behavioural based workbook for cancer patients and e-learning programme for clinicians. A number of studies have also described transplant preparation programmes and reported positive psychological outcomes in patients and their extended network. Davis, Ryninks, Cliss, and Dolby (2013) evaluated the renal transplant preparation programme (RTPP) offered at Bristol Royal Hospital for Children and found that 96% of parents/carers and 100% of children and young people felt that they were given all the information they needed about their transplant. Talking to different members of the team over a number of appointments and settings, as well as building relationships with various professionals offering a person-centred approach, helped patients and their families to prepare for the transplant and the outcomes of their surgery (Davis et al., 2013).


Rolving et al. (2016) used pre-operative cognitive behavioural education to prepare lumbar spinal fusion patients and Grocott, Plumb, Edwards, Fecher-Jones, and Levett (2017) described the use of classroom-based surgery schools to manage patient expectations in relation to the in-hospital surgical journey and improve psychological preparation for surgery. Williamson and Wallace (2012) stated that ‘health professionals are central to the success of any attempt to help patients manage their treatment-related appearance concerns’ (Williamson & Wallace, 2012, p. 431). It is therefore vital that health professionals provide opportunities for patients to raise and openly discuss their concerns about appearance, as well as to be aware of the potential risk factors for poor psychological adjustment to scarring following planned surgery. However, the contribution that health professionals can make to improve patient preparation for surgery is limited by the time available in the care pathway of some specialties between meeting the patient for the first time and the date of surgery (Grocott et al., 2017). Re-designing the pathways to surgery may therefore open opportunities in the future to improve patient care and ultimately improve psychological adjustment to scarring following planned surgery.

### **Models of Adjustment, Disfigurement and Appearance-Related Disorders**

Over the last forty years, a number of models, theories and frameworks have been developed to understand the psychosocial impact of disfigurement and appearance, as well as the maintenance of a number of appearance-related disorders. Rumsey and Harcourt (2004) explored the psychosocial implications of living with a visible difference and identified five models that have been used to explain people’s experiences of disfigurement: Dropkin (1989); Heason (2003); Newell (2000); Kent (2000) and White (2000). However, Rumsey and Harcourt (2004) concluded that the success and utility of the models was limited and required further refinement and testing. Thompson (2012) argued that one model or theory is unlikely to account for the diversity of human experience, or the breadth of activity across body image, disfigurement, attractiveness and identity in appearance-related research. Thompson (2012) highlighted the need for an inclusive and integrative framework to understand disfigurement and this is a view that has been echoed by others (Cash & Pruzinsky, 1990; Grogan, 2016; Newell, 2000; Price, 1990, 1999; Rumsey & Harcourt, 2012). We review the most applicable models next, with a view to informing a theoretical framework for surgical scarring following planned surgery and its consequences.

# **The ARC Framework of Adjustment to Visible Difference (Clarke, Thompson, Jenkinson, Rumsey, & Newell, 2014)**

The Appearance Research Collaboration (ARC) developed a framework to guide their research programme and facilitate the identification of factors contributing to resilience or with the potential to be amenable to change (Clarke et al., 2014). The initial working framework was refined following the analysis and synthesis of the results and the final framework (The ARC Framework of Adjustment to Visible Difference), incorporating the results of the ARC programme can be found in figure 1.1.



*Figure 1.1. The ARC Framework of Adjustment to Visible Difference*

[Reproduced from Clarke et al. (2014) CBT for appearance anxiety: psychosocial interventions for anxiety due to visible difference, p.270. © 2014 John Wiley & Sons, Ltd, with kind permission]

The ARC Framework of Adjustment to Visible Difference incorporates significant aspects of previous models (Cash, 2002; Gilbert, 2002; Kent & Thompson, 2002) with the main findings and trends in appearance research. Within the framework, the process of adjustment to visible difference is made up of three facets: predisposing factors, intervening cognitive processing and outcomes (Clarke et al., 2014). Firstly, predisposing factors are implicated in the development of core beliefs about the self, world and others and include demographic factors such as age and gender, as well as the cause, treatment history and visibility of the site. Secondly, a number of intervening cognitive processes are understood

to process appearance-related information and maintain the overall appearance schema. These include the dispositional style (optimism/pessimism) of the individual, as well as a number of socio-cognitive factors (perceptions of social support/social acceptance, fear of negative evaluations) and appearance-specific cognitions. The final facet is the outcomes of having a visible difference, including appearance-related anxiety, wellbeing, mood, aggression and shame. Further explanation of the framework and findings of the ARC can be found in Clarke et al. (2014).


### **A Cognitive Behavioural Model of Body Dysmorphic Disorder (BDD)**

Alongside the development of models, theories and frameworks to understand the psychosocial impact of disfigurement and appearance more broadly, a number of models have been developed to understand the development and maintenance of specific appearance-related conditions and disorders such as Body Dysmorphic Disorder (BDD). BDD can be understood as ‘a preoccupation with an imagined defect in one’s appearance or, in the case of a slight anomaly, the person’s concern is markedly excessive’ (Veale, 2004, p. 113). The person must also be significantly distressed or handicapped in his or her occupational and social functioning (American Psychiatric Association, 2013). Over the last twenty years, three cognitive behavioural models have been proposed by Wilhelm (Wilhelm, Buhlmann, Hayward, Greenberg, & Dimaite, 2010; Wilhelm, Phillips, & Steketee, 2013), Veale (Veale, 2001; Veale, 2004; Veale et al., 1996) and Neziroglu (Neziroglu, Roberts, & Yaryura-Tobias, 2004) to understand the maintenance of BDD. The cognitive behavioural model of BDD (Veale, 2001; Veale, 2004; Veale et al., 1996) has strong empirical evidence and treatment derived from it has now been tested in two randomised controlled trials (Veale, 2004; Veale et al., 2014).

The model of BDD (Veale, 2004) begins with the trigger of an external representation of an individual’s appearance (e.g. reflection in a mirror or photograph from when they were younger) activating a distorted mental image (figure 1.2). The process of selective and self-focused attention causes the individual to focus on specific aspects of the external representation and leads to heightened awareness of the image and an exaggeration of certain features within it. The individual uses this distorted mental image to construct how they look to others. Assumptions and values relating to appearance and its importance are activated and result in negative appraisal of appearance and processing of self as an aesthetic object. The cognitive behavioural model of BDD predicts that negative appraisals and constant appearance comparisons (either with an ideal internal image or with others) will have a negative feedback loop and increase self-focused attention on the image. It is likely that attention will also interfere with the individual’s ability to process other external



information. Moreover, increases in emotional arousal tend to increase the frequency or severity of negative appraisal of one's body image and further increase self-focused attention. While safety seeking behaviours are often employed to decrease distress or uncertainty, they are counter-productive and increase self-consciousness, preoccupation and negative appraisal (Veale, 2004).



*Figure 1.2.* The cognitive behavioural model of BDD

[Reproduced from Veale (2004) *Advances in a cognitive behavioural model of body dysmorphic disorder*, with kind permission]

A number of studies have further investigated specific aspects of the BDD model, as outlined in Baldock and Veale (2017), including: selective attention and heightened awareness (Anson, Veale, & de Silva, 2012; Clerkin & Teachman, 2008; Kollei & Martin, 2014); distorted mental image (Osman, Cooper, Hackmann, & Veale, 2004; Willson, Veale, & Freeston, 2016); mirror gazing (Baldock, Anson, & Veale, 2012; Windheim, Veale, & Anson, 2011); and self as aesthetic object (Lambrou, Veale, & Wilson, 2011, 2012; Veale, Kinderman, Riley, & Lambrou, 2003).

## **A New Understanding of Psychological Adjustment to Scarring following Planned Surgery**

Although a number of models, theories and frameworks have been developed to understand the psychosocial impact of disfigurement and the maintenance of a number of appearance-related disorders, they have largely focused on when the disfigurement is already present and do not necessarily aid understanding of adjustment to planned surgery. Notably, the circumstance by which a scar is acquired and how the acquisition is managed are also important and will play a role in an individual's adjustment. Within the field of surgical scarring, it is the treatment that causes the scarring in the first place. It may therefore be helpful to intervene prior to surgery to prevent some of the difficulties described earlier in this review. Rather than propose a completely separate model to understand disfigurement, our new model (figure 1.3) builds on those previously described in the literature to better understand psychological adjustment to scarring following planned surgery. The model draws on previous research to focus on areas where health professionals might better support their patients by providing specific and targeted preparation and treatment in future. It promotes early preparation and intervention prior to surgery, with the aim of improving adjustment and psychological outcomes in the long-term, and helps identify promising research areas.

### **A Model of Psychological Adjustment to Scarring following Planned Surgery**

The model presented here is composed of two main components: developmental and triggering aspects which contribute to meaning, and maintaining factors which arise from and can in turn influence those meanings. These are predominantly drawn from the literature reviewed above within a cognitive-behavioural theoretical framework.

#### **Before the Operation/Surgery**

Taking developmental and triggering factors first. A number of pre-existing factors (as outlined previously by Clarke et al. (2014)) may influence an individual's anticipated experience of their surgery and their reactions post-operatively. Katz et al. (2003) reported that women experienced lower life happiness and greater depressive symptoms following treatment for head and neck cancer than men, as well as more appearance dissatisfaction and concern (Khafif et al., 2007; Liu, 2008). Nissen et al. (2001) found older women tend to report more positive body image following breast cancer surgery than younger women and Hood (2010) reported appearance concerns were less important in older women. Gender and age may therefore play a role in adjustment, although evidence from other studies suggests that the role of these factors may not be so clear cut.




Figure 1.3. A Model of Psychological Adjustment to Scarring following Planned Surgery.

This model is meant to summarise the full range of adjustment to scarring following planned surgery, from good to problematic. The upper part of the figure shows influences which have been noted to mediate or moderate the meaning attached to scarring; the lower section indicates processes which, if the meaning is particularly negative, may maintain psychological problems occurring as a consequence of scarring.

Societal and cultural influences (including the media) are also important when considering adjustment to scarring. Beliefs that society places a high value on attractiveness (Tebble et al., 2004; Versnel et al., 2012), as well as social connotations of femininity, sexuality and motherhood of the chest region (Figueiredo et al., 2004; Khan et al., 2000) may feed into an individual's anticipated experience and resulting perceived impact of surgical scarring. Sociocultural influences therefore need to be considered as part of preparing individuals for surgery (whilst simultaneously bearing in mind the ultimate need to change attitudes about appearance at a societal level, rather than simply supporting individuals in this regard). The developmental stage within which the illness is acquired and the developmental stage in which the operation occurred or scar is acquired are also important and will affect an individual's adjustment to scarring following surgery. Kantoch et al. (2006) reported that scars acquired in adulthood exerted less impact on self-esteem and self-confidence than those acquired in childhood or adolescence.

Siegel (2010) reported that the foundation of our sense of self is built upon the intimate exchanges we have as babies. Where the exchanges have been positive, individuals develop secure attachments. In contrast, where the exchanges have been less positive, individuals may develop a variety of insecure attachment styles. A positive sense of self and self-image are developed through secure attachments with parents and caregivers (Sroufe, 1995, 1997) and these individuals score highly on resilience, adaptability and self-esteem. Factors such as bullying in childhood can contribute to an individual's increased risk of developing adverse physical health, emotional and psychological outcomes in childhood (Reijntjes, Kamphuis, Prinzie, & Telch, 2010), as well as a poor sense of self and increased risk of poor health, wealth and social-relationship outcomes in adulthood (Wolke, Copeland, Angold, & Costello, 2013). Individuals with a less positive sense of self and self-image may therefore be more likely to have poorer adjustment to scarring following surgery.

In addition, temperament (broadly defined by Derryberry and Rothbart (1988) as an emotional and behavioural style appearing in early life that is consistent across time) and more specifically negative affectivity (a tendency towards discomfort, fear, anger, sadness, and low soothability) have been shown to be linked to anxiety and depression, as well as maladaptive coping styles (Miller et al., 2009). Since temperament is less malleable than coping style, it is likely that certain temperamental traits may prevent some individuals from using effective coping strategies following surgery and consequently have poorer outcomes. As with previous models of adjustment and disfigurement, pre-existing factors influence the meaning that the individual makes of their changed appearance (surgical scar). Individuals with a more positive sense of self and self-image and who attach less meaning to their

appearance and body image may adjust better following surgical scarring. Pre-operative factors also feed into an individual's anticipated experience, with evidence suggesting that a lack of preparation may result in individuals being poorly-prepared to live with their scar (King et al., 2009). Psychological and medical preparation (e.g. anticipated medical outcomes and patient journey, length of recovery) are therefore critical to individuals being prepared for all eventualities of the surgery.

Family beliefs and behaviour are also involved in adjustment, as particular parenting styles and family dynamics will affect how the family and individual prepare for their operation and experience illness. It is therefore important to ensure that the wider family is also prepared and supported, rather than only focusing attention on the individual undergoing the surgery. Whilst some families may encourage discussions about their feelings around the operation and illness and address any that may be considered problematic, others may avoid discussions completely or only talk at the last minute. Evidence suggests that sharing and talking about one's thoughts, emotions and experiences in relation to health results in more positive adjustment (Mallinger, Griggs, & Shields, 2006). Manne, Pape, Taylor, and Dougherty (1999) found that patients undergoing active treatment for breast cancer who perceived that their spouses did not want to talk about cancer-related issues experienced greater psychological distress. Finally, pre-operative and pre-existing factors influence each other in a bidirectional way, as attachment style, temperament and other demographic factors will affect how the individual and their family prepare for the operation.

### **After the Operation/Surgery**

The anticipated experience is important when considering psychological adjustment to scarring as it influences the individual's resulting perceived impact of the operation and post-op reaction and meaning. Although surgical scarring is the result of an operation that broadly aims to improve an individual's health and/or functioning, the operation and scar may not occur as planned. Whilst the eventual medical outcomes of surgery contribute to the meaning that the individual makes of their changed appearance, a number of dynamic factors (short term/immediate medical outcomes) also play a role. For example, the circumstances around the operation itself are important as they may mean a change in the anticipated pathway of treatment. Complications and a long stay in intensive care may take individuals by surprise and make them question beliefs about mortality and their patient journey. Anecdotal observations suggest many individuals focus upon their desired positive outcome (improved health and/or functioning), without consideration of the process (surgery) they must go through in order to obtain their positive outcome in the first place.

Although a focus on outcome may be an effective coping strategy for some, for individuals where there are complications (the scar is worse than anticipated or surgery does not produce the desired or anticipated positive outcome) psychological adjustment to surgical scarring may be poor. Individuals' beliefs about the level of pain they will experience, their likelihood of dying in surgery and loss of control all contribute to their anticipated experience, with factors such as increased pain negatively influencing the meaning they make of their surgery. Therefore, rather than the scar representing a symbol of their positive outcome, it may now represent disappointment and anger at an unsuccessful outcome or complications they had to endure.

In addition, the reactions of others (both perceived and actual) impact upon the meaning that an individual makes of their surgery and changed appearance. Partridge (1990) highlighted the varied reactions of family and friends to facial disfigurements, with some committed to supporting their loved one and others avoiding or rejecting the individual who had been disfigured. Some patients with head and neck cancer have reported feeling stigmatised, with difficulty making friends and securing employment following treatment (Kent, 2000). Piot-Ziegler et al. (2010) found women frequently reported difficulties with intimacy following surgery and attempted to conceal their scars for fear of stigma or shocking others. Regardless of the cause of disfigurement (acquired versus congenital), the most frequently reported difficulties relate to encounters with strangers, meeting new people and making new friends (Robinson, 1997). It is therefore important to consider the reactions of others as part of preparation for surgery, including the possibility of a self-fulfilling prophecy where individuals may anticipate negative reactions and therefore behave in a shy or defensive manner, in turn inviting negativity from others.

Crucially, the meaning that an individual attaches to the process they have been through and its outcomes results in a number of behavioural changes and processes that have been well described previously and determine post-operative adjustment, including social and self-ideal comparison, safety-seeking and other motivated behaviours (e.g. attempts to conceal scars, Piot-Ziegler et al. (2010)), rumination, selective attention and mood (e.g. anxiety and depression, Semple et al. (2004)). Where the meaning is particularly negative, this will be maintained to the extent that these responses hold it in place or increase it in terms of negativity, in the way seen in persistent health anxiety (Salkovskis & Warwick, 1986) and BDD (Veale, 2004).

## **Summary**

Over the last fifty years, a number of models, theories and frameworks have been developed to understand the psychosocial impact of disfigurement and the maintenance of a number of appearance-related disorders. They have largely focused on understanding adjustment when the disfigurement is already present. However, the circumstance by which a scar is acquired and how the acquisition is managed are also important and will play a role in an individual's adjustment. This review highlights the importance of shifting the focus in medicine from 'functioning body parts' to positive psychological adjustment and wellbeing following surgery. The management of scars presents a challenge for health professionals, as scarring can greatly impact upon quality of life, with proneness to shame, body image problems and lack of self-esteem (Bock et al., 2006; França et al., 2013). Within the field of surgical scarring, it is the treatment that causes the scarring in the first place. Psychological preparation for surgery may help to reduce this challenge and patient information and preparedness have long been considered key elements of enhanced recovery pathways (Grocott et al., 2012).

Psychologists working in health settings have been asking for a model to formulate and improve psychological outcomes in their patients. Rather than propose a completely separate model to understand disfigurement, our model builds on those previously described in the literature (Clarke et al., 2014; Veale, 2004) to better understand psychological adjustment to scarring following planned surgery. The model draws on previous research to focus on areas where health professionals might better support their patients by providing specific and targeted preparation (preventatively focused) and treatment where scar related distress becomes persistent. It promotes early preparation and intervention prior to surgery, with the aim of preventing some of the difficulties described earlier in this review. This should have the benefit of improving adjustment to surgical scarring and psychological outcomes in the long-term, and helps to identify promising areas of research. Moreover, elements of our model may be useful in understanding adjustment in other appearance-related conditions where context and preparation might be key.

## **Clinical Implications**

The literature in this review highlights that it is the patient's subjective opinion of their scar that counts when considering adjustment. Health professionals may objectively see the scar in a 'positive' light and make assumptions about how the patient will respond; however, it is important for health professionals to consider the patient's perception of their scar, how they may respond to it (e.g. conceal) and how it may be perceived by others in

order to tailor their approach to patients accordingly. Evidence from the BDD literature highlights the challenges of helping patients to talk about their appearance concerns (Marques, Weingarden, LeBlanc, & Wilhelm, 2011). Health professionals therefore need to feel confident in raising these issues with their patients and may benefit from training in how to ask the right questions, and not to take answers at face value. It is also vital that health professionals are aware of the potential risk factors for poor psychological adjustment to scarring following planned surgery, and that they provide opportunities for patients to raise and openly discuss their concerns about appearance and medical outcomes. Our model provides a framework for multidisciplinary teams to do this. Helping patients to understand and process the stories of their scars (e.g. successes and complications) may also help them to create meaning (e.g. reminder of strength or bravery vs. sign of weakness). Health professionals can therefore model positive story-telling to patients by talking through the more difficult aspects of the surgery and helping patients and their families to re-frame and feel more able to tell their stories themselves (see Davis et al., in preparation, for further discussion). Where concerns about scarring remain, health professionals may find it useful to refer patients for psychological support or signpost to appearance concern websites and charities (e.g. Changing Faces and YP Face IT). Care pathways may need to be redesigned to create space for these conversations to occur and new pathways could be developed for those where surgical procedures are expected to result in scarring. Such a care pathway could include the development of new tools for surgical preparation and a post-surgical stepped-care approach to psychological reactions. Self-help and psychoeducation material (drawing on the understanding summarised in this paper) can be provided as part of routine care. More focused self-help materials can be used post-operatively for individuals experiencing difficulties in adjustment and where problems persist and substantially impact on quality of life, treatment drawing on the model defined here could be offered. However, making changes to existing care pathways is contingent upon service managers and commissioners seeing the value in working in this way. It is hoped that psychologists working in health settings can use this model to provide teaching and training to their multidisciplinary colleagues to increase confidence in holding these conversations with patients and their families in the future.



## References

- Alfano, C. M., & Rowland, J. H. (2006). Recovery issues in cancer survivorship: a new challenge for supportive care. *The Cancer Journal*, 12(5), 432-443.
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders: DSM-5* (5th ed.). Washington, DC.
- Anson, M., Veale, D., & de Silva, P. (2012). Social-evaluative versus self-evaluative appearance concerns in Body Dysmorphic Disorder. *Behaviour research and therapy*, 50(12), 753-760.
- Baldock, E., Anson, M., & Veale, D. (2012). The stopping criteria for mirror-gazing in body dysmorphic disorder. *British Journal of Clinical Psychology*, 51(3), 323-344.
- Baldock, E., & Veale, D. (2017). The self as an aesthetic object: a model of cognitive processing in Body Dysmorphic Disorder. In K. A. Phillips (Ed.), *Body Dysmorphic Disorder: Advances in Research and Clinical Practice* (pp. 299-310): Oxford University Press.
- Bayat, A., McGrouther, D., & Ferguson, M. (2003). Skin scarring. *BMJ: British Medical Journal*, 326(7380), 88.
- Bernstein, N. (1982). Psychosocial results of burns. The damaged self-image. *Clinics in plastic surgery*, 9(3), 337-346.
- Beuf, A. H. (1990). *Beauty is the beast: Appearance-impaired children in America*. Philadelphia: University of Pennsylvania Press.
- Bock, O., Schmid-Ott, G., Malewski, P., & Mrowietz, U. (2006). Quality of life of patients with keloid and hypertrophic scarring. *Archives of dermatological research*, 297(10), 433-438.
- Brown, B., McKenna, S., Siddhi, K., McGrouther, D., & Bayat, A. (2008). The hidden cost of skin scars: quality of life after skin scarring. *Journal of Plastic, Reconstructive & Aesthetic Surgery*, 61(9), 1049-1058.
- Brown, B., Moss, T., McGrouther, D., & Bayat, A. (2010). Skin scar preconceptions must be challenged: importance of self-perception in skin scarring. *Journal of Plastic, Reconstructive & Aesthetic Surgery*, 63(6), 1022-1029.

- Bull, R., & Rumsey, N. (2012). *The social psychology of facial appearance*. New York: Springer-Verlag.
- Cash, T. F. (2002). Cognitive-behavioral perspectives on body image. In T. F. Cash & T. Pruzinsky (Eds.), *Body image: A handbook of theory, research, and clinical practice* (pp. 38-46). London: The Guildford Press.
- Cash, T. F., & Pruzinsky, T. (1990). *Body images: Development, deviance, and change*: Guilford Press.
- Chacon, A., França, K., Ledon, J., & Nouri, K. (2013). Scars and Scarring: Causes, Types and Treatment Options *Scars and Scarring: Causes, Types and Treatment Options* (pp. 1-24): Nova Science Publishers, Inc.
- Clarke, A., Thompson, A. R., Jenkinson, E., Rumsey, N., & Newell, R. (2014). *CBT for appearance anxiety: psychosocial interventions for anxiety due to visible difference*: John Wiley & Sons.
- Clerkin, E. M., & Teachman, B. A. (2008). Perceptual and cognitive biases in individuals with body dysmorphic disorder symptoms. *Cognition and Emotion*, 22(7), 1327-1339.
- Crossland, D., Jackson, S., Lyall, R., Hamilton, J., Hasan, A., Burn, J., & O'Sullivan, J. (2005). Patient attitudes to sternotomy and thoracotomy scars. *The thoracic and cardiovascular surgeon*, 53(02), 93-95.
- Davis, C., Ellison, N., Sheils, L., King, G., Veale, D., Garratt, V., . . . Tulloh, R. (in preparation). Children living with heart surgery scars: War wounded but armed with a story?
- Davis, C., Ryninks, K., Cliss, A., & Dolby, S. (2013). A partnership approach to transplant preparation: Involving children, young people and their families. *British Journal of Renal Medicine*, 18(4), 26-29.
- Derryberry, D., & Rothbart, M. K. (1988). Arousal, affect, and attention as components of temperament. *Journal of personality and social psychology*, 55(6), 958.
- Dropkin, M. J. (1989). *Coping with disfigurement and dysfunction after head and neck cancer surgery: a conceptual framework*. Paper presented at the Seminars in Oncology Nursing.

- Figueiredo, M. I., Cullen, J., Hwang, Y.-T., Rowland, J. H., & Mandelblatt, J. S. (2004). Breast cancer treatment in older women: does getting what you want improve your long-term body image and mental health? *Journal of Clinical Oncology*, 22(19), 4002-4009.
- Fingeret, M. C., Yuan, Y., Urbauer, D., Weston, J., Nipomnick, S., & Weber, R. (2012). The nature and extent of body image concerns among surgically treated patients with head and neck cancer. *Psycho-Oncology*, 21(8), 836-844.
- Fobair, P., Stewart, S. L., Chang, S., D'onofrio, C., Banks, P. J., & Bloom, J. R. (2006). Body image and sexual problems in young women with breast cancer. *Psycho-Oncology*, 15(7), 579-594.
- França, K., Chacon, A., Ledon, J., Savas, J., & Nouri, K. (2013). Psychological Impact of Scars *Scars and Scarring: Causes, Types and Treatment Options* (pp. 25-38): Nova Science Publishers, Inc.
- Gamba, A., Romano, M., Grosso, L. M., Tamburini, M., Cantú, G., Molinari, R., & Ventafridda, V. (1992). Psychosocial adjustment of patients surgically treated for head and neck cancer. *Head & neck*, 14(3), 218-223.
- Gaskell, S. (2010). Evidence-based guidelines for the management of invasive and/or distressing procedures with children. *Brit Psychol Soc*, 1-44.
- Gauglitz, G. G., Korting, H. C., Pavicic, T., Ruzicka, T., & Jeschke, M. G. (2011). Hypertrophic scarring and keloids: pathomechanisms and current and emerging treatment strategies. *Mol Med*, 17(1-2), 113-125.
- Geyer, S., Norozi, K., Zoege, M., Kempa, A., Buchhorn, R., & Wessel, A. (2006). Psychological symptoms in patients after surgery for congenital cardiac disease. *Cardiology in the Young*, 16(06), 540-548.
- Gilbert, P. (2002). Body shame: A biopsychosocial conceptualisation and overview, with treatment implications. In P. Gilbert & J. Miles (Eds.), *Body shame: Conceptualisation, research and treatment* (pp. 3-54). Hove: Brunner-Routledge.
- Grocott, M. P., Martin, D. S., & Mythen, M. G. (2012). Enhanced recovery pathways as a way to reduce surgical morbidity. *Current opinion in critical care*, 18(4), 385-392.

- Grocott, M. P., Plumb, J. O., Edwards, M., Fecher-Jones, I., & Levett, D. Z. (2017). Re-designing the pathway to surgery: better care and added value. *Perioperative Medicine*, 6(1), 9.
- Grogan, S. (2016). *Body image: Understanding body dissatisfaction in men, women and children*: Routledge.
- Heason, S. L. (2003). *The development of a model of disfigurement: The process of living with vitiligo*. University of Sheffield.
- Hood, C. (2010). Project to improve care for people coping with changes in body image: A study to understand the experiences of patients coping with life during and after treatment resulted in the development of a toolkit to enable healthcare professionals to provide individualised care and support. Caroline Hood reports. *Cancer Nursing Practice*, 9(2), 26-32.
- Kantoch, M. J., Eustace, J., Collins-Nakai, R. L., Taylor, D. A., Boisvert, J. A., & Lysak, P. S. (2006). The significance of cardiac surgery scars in adult patients with congenital heart disease. *Kardiologia polska*, 64(1), 51-56.
- Katz, M. R., Irish, J. C., Devins, G. M., Rodin, G. M., & Gullane, P. J. (2003). Psychosocial adjustment in head and neck cancer: the impact of disfigurement, gender and social support. *Head & neck*, 25(2), 103-112.
- Kent, G. (2000). Understanding the experiences of people with disfigurements: an integration of four models of social and psychological functioning. *Psychology, health & medicine*, 5(2), 117-129.
- Kent, G., & Thompson, A. R. (2002). Models of disfigurement: Implications for treatment. In P. Gilbert & J. Miles (Eds.), *Understanding body shame* (pp. 106-116). Hove: Brunner-Routledge.
- Khafif, A., Posen, J., Yagil, Y., Beiser, M., Gil, Z., Ben-Yosef, R., . . . Fliss, D. M. (2007). Quality of life in patients older than 75 years following major head and neck surgery. *Head & neck*, 29(10), 932-939.
- Khan, M., Sehgal, A., Mitra, A., Agarwal, P., Lal, P., & Malik, V. (2000). Psycho-behavioural impact of mastectomy. *Journal of the Indian Academy of Applied Psychology*, 26(1-2), 65-71.

- King, K. M., McFetridge-Durdle, J., LeBlanc, P., Anzarut, A., & Tsuyuki, R. T. (2009). A descriptive examination of the impact of sternal scar formation in women. *European Journal of Cardiovascular Nursing*, 8(2), 112-118.
- Klinge, K., Chamberlain, D. J., Redden, M., & King, L. (2009). Psychological adjustments made by postburn injury patients: an integrative literature review. *Journal of advanced nursing*, 65(11), 2274-2292.
- Kollei, I., & Martin, A. (2014). Body-related cognitions, affect and post-event processing in body dysmorphic disorder. *Journal of behavior therapy and experimental psychiatry*, 45(1), 144-151.
- Lambrou, C., Veale, D., & Wilson, G. (2011). The role of aesthetic sensitivity in body dysmorphic disorder. *Journal of abnormal psychology*, 120(2), 443-453.
- Lambrou, C., Veale, D., & Wilson, G. (2012). Appearance concerns comparisons among persons with body dysmorphic disorder and nonclinical controls with and without aesthetic training. *Body image*, 9(1), 86-92.
- Lawrence, J. W., Mason, S. T., Schomer, K., & Klein, M. B. (2012). Epidemiology and impact of scarring after burn injury: a systematic review of the literature. *Journal of Burn Care & Research*, 33(1), 136-146.
- Liu, H. E. (2008). Changes of satisfaction with appearance and working status for head and neck tumour patients. *Journal of clinical nursing*, 17(14), 1930-1938.
- Lupton, J. R., & Alster, T. S. (2002). Laser scar revision. *Dermatologic clinics*, 20(1), 55-65.
- MacGregor, F. C. (1970). Social and psychological implications of dentofacial disfigurement. *The Angle Orthodontist*, 40(3), 231-233.
- Mallinger, J., Griggs, J., & Shields, C. (2006). Family communication and mental health after breast cancer. *European Journal of Cancer Care*, 15(4), 355-361.
- Manne, S. L., Pape, S. J., Taylor, K. L., & Dougherty, J. (1999). Spouse support, coping, and mood among individuals with cancer. *Annals of Behavioral Medicine*, 21(2), 111-121.

- Marques, L., Weingarden, H. M., LeBlanc, N. J., & Wilhelm, S. (2011). Treatment utilization and barriers to treatment engagement among people with body dysmorphic symptoms. *Journal of psychosomatic research*, 70(3), 286-293.
- Masi, G., & Brovedani, P. (1999). Adolescents with congenital heart disease: psychopathological implications. *Adolescence*, 34(133), 185.
- Miller, K. S., Vannatta, K., Compas, B. E., Vasey, M., McGoron, K. D., Salley, C. G., & Gerhardt, C. A. (2009). The role of coping and temperament in the adjustment of children with cancer. *Journal of pediatric psychology*, 34(10), 1135-1143.
- Moss, T. P. (2005). The relationships between objective and subjective ratings of disfigurement severity, and psychological adjustment. *Body image*, 2(2), 151-159.
- Newell, R. (2000). *Body image and disfigurement care*. London: Routledge.
- Neziroglu, F., Roberts, M., & Yaryura-Tobias, J. A. (2004). A behavioral model for body dysmorphic disorder. *Psychiatric Annals*, 34(12), 915-920.
- Nissen, M. J., Swenson, K. K., Ritz, L. J., Farrell, J. B., Sladek, M. L., & Lally, R. M. (2001). Quality of life after breast carcinoma surgery. *Cancer*, 91(7), 1238-1246.
- Osman, S., Cooper, M., Hackmann, A., & Veale, D. (2004). Spontaneously occurring images and early memories in people with body dysmorphic disorder. *Memory*, 12(4), 428-436.
- Papaspyros, S., Patel, R., Polyzois, K., Javagula, K., & Jeffrey, R. (2011). Median sternotomy scars: formation and impact on patient quality of life. *British Journal of Cardiac Nursing*, 6(11), 531-540.
- Philipp, C. M., Scharschmidt, D., & Berlien, H. P. (2008). Laser treatment of scars and keloids—How we do it. *Medical Laser Application*, 23(2), 79-86.
- Piff, C. (1998). Body image: a patient's perspective. *The British journal of theatre nursing: NATNews: the official journal of the National Association of Theatre Nurses*, 8(1), 13-14.
- Piot-Ziegler, C., Sassi, M. L., Raffoul, W., & Delaloye, J. F. (2010). Mastectomy, body deconstruction, and impact on identity: a qualitative study. *British journal of health psychology*, 15(3), 479-510.

- Price, B. (1990). A model for body-image care. *Journal of advanced nursing*, 15(5), 585-593.
- Price, B. (1999). *Altered body image Nursing Times Monographs No. 29*. London: NT Books.
- Rajipura, A. (2002). *The National Burn Repository Report 2002*. Retrieved from <http://www.ameriburn.org>
- Reijntjes, A., Kamphuis, J. H., Prinzie, P., & Telch, M. J. (2010). Peer victimization and internalizing problems in children: A meta-analysis of longitudinal studies. *Child abuse & neglect*, 34(4), 244-252.
- Robinson, E. (1997). Psychological research on visible differences in adults. In R. Lansdown, N. Rumsey, E. Bradbury, A. Carr, & J. Partridge (Eds.), *Visibly different: Coping with disfigurement* (pp. 102-111). Oxford: Butterworth-Heinemann.
- Rolving, N., Sogaard, R., Nielsen, C. V., Christensen, F. B., Bünger, C., & Oestergaard, L. G. (2016). Preoperative cognitive-behavioral patient education versus standard care for lumbar spinal fusion patients: economic evaluation alongside a randomized controlled trial. *Spine*, 41(1), 18-25.
- Royal College of Surgeons. (2016). *Surgery and the NHS in numbers*.
- Rumsey, N., Clarke, A., White, P., Wyn-Williams, M., & Garlick, W. (2004). Altered body image: appearance-related concerns of people with visible disfigurement. *Journal of advanced nursing*, 48(5), 443-453.
- Rumsey, N., & Harcourt, D. (2004). Body image and disfigurement: issues and interventions. *Body image*, 1(1), 83-97.
- Rumsey, N., & Harcourt, D. (2012). *The Oxford Handbook of the Psychology of Appearance*. Oxford: Oxford University Press.
- Salkovskis, P. M., & Warwick, H. M. (1986). Morbid preoccupations, health anxiety and reassurance: a cognitive-behavioural approach to hypochondriasis. *Behaviour research and therapy*, 24(5), 597-602.

- Sarwer, D., Whitaker, L., & Bartlett, S. (2001). *Psychological functioning of adolescents born with craniofacial anomalies*. Paper presented at the 9th International Congress of the International Society of Craniofacial Surgery. Craniofacial Surgery.
- Semple, C. J., Sullivan, K., Dunwoody, L., & Kernohan, W. G. (2004). Psychosocial interventions for patients with head and neck cancer: past, present, and future. *Cancer nursing*, 27(6), 434-441.
- Shakespeare, V. (1998). Effect of small burn injury on physical, social and psychological health at 3–4 months after discharge. *Burns*, 24(8), 739-744.
- Siegel, D. J. (2010). *Mindsight: The New Science of Personal Transformation*: Random House.
- Sroufe, A. L. (1995). *Emotional Development*. Cambridge: Cambridge University Press.
- Sroufe, A. L. (1997). Psychopathology as an outcome of development. *Development and psychopathology*, 9(2), 251-268.
- Tebble, N. J., Thomas, D. W., & Price, P. (2004). Anxiety and self-consciousness in patients with minor facial lacerations. *Journal of advanced nursing*, 47(4), 417-426.
- Tedstone, J. E., & Tarrier, N. (1997). An investigation of the prevalence of psychological morbidity in burn-injured patients. *Burns*, 23(7), 550-554.
- Thomas, C. S. (1990). Stress and facial appearance. *Stress Medicine*, 6(4), 299-304.
- Thompson, A. R. (2012). Researching Appearance: Models, Theories, and Frameworks. In N. R. a. D. Harcourt (Ed.), *The Oxford Handbook of the Psychology of Appearance* (pp. 91-109). Oxford: Oxford University Press.
- Thompson, A. R., Kent, G., & Smith, J. A. (2002). Living with vitiligo: dealing with difference. *British journal of health psychology*, 7(2), 213-225.
- Van Loey, N. E., & Van Son, M. J. (2003). Psychopathology and psychological problems in patients with burn scars. *American journal of clinical dermatology*, 4(4), 245-272.
- Veale, D. (2001). Cognitive-behavioural therapy for body dysmorphic disorder. *Advances in psychiatric treatment*, 7(2), 125-132.
- Veale, D. (2004). Advances in a cognitive behavioural model of body dysmorphic disorder. *Body image*, 1(1), 113-125.



- Veale, D., Anson, M., Miles, S., Pieta, M., Costa, A., & Ellison, N. (2014). Efficacy of cognitive behaviour therapy versus anxiety management for body dysmorphic disorder: a randomised controlled trial. *Psychotherapy and psychosomatics*, 83(6), 341-353.
- Veale, D., Gournay, K., Dryden, W., Boocock, A., Shah, F., Willson, R., & Walburn, J. (1996). Body dysmorphic disorder: a cognitive behavioural model and pilot randomised controlled trial. *Behaviour research and therapy*, 34(9), 717-729.
- Veale, D., Kinderman, P., Riley, S., & Lambrou, C. (2003). Self-discrepancy in body dysmorphic disorder. *British Journal of Clinical Psychology*, 42(2), 157-169.
- Versnel, S. L., Plomp, R. G., Passchier, J., Duivenvoorden, H. J., & Mathijssen, I. M. (2012). Long-term psychological functioning of adults with severe congenital facial disfigurement. *Plastic and reconstructive surgery*, 129(1), 110-117.
- White, C. A. (2000). Body image dimensions and cancer: a heuristic cognitive behavioural model. *Psycho-Oncology*, 9(3), 183-192.
- Wilhelm, S., Buhlmann, U., Hayward, L. C., Greenberg, J. L., & Dimaite, R. (2010). A cognitive-behavioral treatment approach for body dysmorphic disorder. *Cognitive and Behavioral Practice*, 17(3), 241-247.
- Wilhelm, S., Phillips, K. A., & Steketee, G. (2013). *Cognitive-Behavioural Therapy for Body Dysmorphic Disorder*. New York: The Guildford Press.
- Williamson, H., & Wallace, M. (2012). When Treatment Affects Appearance. In N. Rumsey & D. Harcourt (Eds.), *Oxford Handbook of the Psychology of Appearance*: Oxford University Press.
- Willson, R., Veale, D., & Freeston, M. (2016). Imagery rescripting for body dysmorphic disorder: A multiple-baseline single-case experimental design. *Behavior Therapy*, 47(2), 248-261.
- Windheim, K., Veale, D., & Anson, M. (2011). Mirror gazing in body dysmorphic disorder and healthy controls: Effects of duration of gazing. *Behaviour research and therapy*, 49(9), 555-564.
- Wolke, D., Copeland, W. E., Angold, A., & Costello, E. J. (2013). Impact of bullying in childhood on adult health, wealth, crime, and social outcomes. *Psychological science*, 24(10), 1958-1970.

Service Improvement Project

**Volunteers' experiences of helping hoarders and hoarders' experiences of being helped**

Kirsty Ryninks

Doctorate in Clinical Psychology

Department of Psychology, University of Bath, Claverton Down, Bath, BA2 7AY

Email: K.E.Ryninks@bath.ac.uk

Word count: 5189

May 2018

**Internal/academic supervisors:** Dr James Gregory and Dr Vuokko Wallace;

Doctoral Programme in Clinical Psychology

Department of Psychology, University of Bath, Claverton Down, Bath, BA2 7AY

Email: J.D.Gregory@bath.ac.uk and V.Wallace@bath.ac.uk

**Proposed Journal:** British Journal of Clinical Psychology

Please see Appendix IV for instructions to authors



## **Introduction**

Hoarding disorder is characterised “by persistent difficulty discarding or parting with possessions” and “causes clinically significant distress or impairment in social, occupational and other important areas of functioning” (American Psychiatric Association, 2013). This difficulty results in the accumulation of possessions and causes living areas to become cluttered and their proposed use significantly compromised. Hoarding has been reported to result in economic, familial and social burden (Frost, Steketee, & Williams, 2000), with the severely restricted use of living areas, along with feelings of shame and embarrassment often creating a barrier to others accessing the home (Coles, Frost, Heimberg, & Steketee, 2003).

Studies have repeatedly reported evidence that symptoms of hoarding emerge in mid-to-late childhood (Frost & Gross, 1993; Grisham, Frost, Steketee, Kim, & Hood, 2006), with symptom severity increasing with every decade of life (Ayers, Saxena, Golshan, & Wetherell, 2010). Eckfield and Wallhagen (2013) reported significantly higher rates of hoarding in adults over 55 years old and according to service providers, hoarding behaviour posed a serious physical threat in 81% of their elderly clients (Kim, Steketee & Frost, 2001). Hoarding has been found to increase the risk of falls, fire hazards, food contamination, social isolation and medication mismanagement (Ayers et al., 2010); hoarding can therefore pose serious consequences for older adults. Although there is high comorbidity with hoarding and other mental health conditions (Mataix-Cols et al., 2010), Pertusa et al. (2010) reported that there is an unwillingness amongst hoarders to acknowledge their problem as a mental health difficulty and to access help from services. Agencies who are responsible for supporting adults in later life consequently provide a key route through which to identify and work with hoarding difficulties (Frost et al., 2000).

A diverse range of professionals are involved in working with hoarders, including social services, medical, fire, environmental and mental health professionals (Slatter, 2007). Social service agencies have tended to employ a fragmented treatment approach, with occasional interventions often resulting in relapse (Damecour & Charron, 1998). The Chartered Institute of Environmental Health (2009) describes involvement of environmental health in cases related to hoarding when complaints have been made by neighbours or there are concerns regarding risk to physical health. When regulations regarding sanitation, antisocial behaviour or health are breached, clearance is mandatory, but evidence suggests that clearance interventions have poor long-term prognosis due to the absence of insight/behaviour change (Chartered Institute of Environmental Health, 2009).

Studies have repeatedly reported modest successes and high rates of drop out when working with this client group (Steketee & Frost, 2007; Turner, Steketee, & Nauth, 2010). Ayers, Wetherell, Golshan, and Saxena (2011) recruited twelve adults aged over 65 years to examine the impact of a manualised cognitive-behavioural therapy (CBT) protocol for compulsive hoarding. Although the findings revealed statistically significant changes to depression scores and hoarding severity, only three adults were identified as treatment responders and at follow-up their progress had not been maintained. Ayers, Bratiotis, Saxena, and Wetherell (2012) used a qualitative approach to explore therapist and patient perspectives of their specialised CBT protocol. The therapeutic relationship, exposure exercises and home visits were reported by patients to be most helpful whereas cognitive-behavioural formulations and techniques were not helpful. Therapists considered session and homework compliance to be critical to treatment outcomes.

Compulsive hoarding is widely thought of as a difficult problem to help with (Tolin, Frost, & Steketee, 2012) and evidence suggests that supporting elderly clients who hoard represents a significant challenge to services (Turner et al., 2010). Most studies to date have focused on understanding the experiences of professionals and public sector workers with people who hoard. Negative and judgmental attitudes, as well as burnout, frustration and helplessness have also been reported (Frost, Tolin, & Maltby, 2010). More recently, Holden, Kellett, Davies, and Scott (2016) explored the experience of professionals (e.g. social care, mental health, environmental health, fire service) working with clients that hoard. They found that professionals experience and approach their work in discrete and dissimilar ways, identifying three distinct clusters of professionals: therapeutic and client-focused, shocked and frustrated, and pragmatic and task focused.

Given the outcome data and reported difficulty of working with hoarders, it is important to understand the factors that act as barriers and facilitators in supporting people with compulsive hoarding. Service constraints and increasingly pressurised financial budgets are causing services to seek new ways to support their clients. Evidence suggests that support from non-professionals may be helpful in enhancing clients' physical and mental wellbeing (Taft et al., 2011; Uhm et al., 2016). The present study aimed to explore the experiences of compulsive hoarders and volunteer helpers within the context of a UK-based charity providing support to older adults with hoarding difficulties.

## **Method**

### **Participants and Recruitment**

Participants were recruited through a UK-based charity that provides advice and assistance to enable older and disabled people to continue living independently. The charity's 'Making Space' project developed with the aim of volunteers providing practical and emotional support to adults with hoarding tendencies. A purposive sample of eleven participants from the Making Space project participated in the study. Four clients (two males and two females), who self-identified as having a problem accumulating numerous possessions, were interviewed in their own homes. Seven volunteers (two males and five females) were interviewed at the charity's office base. Ethical approval was granted by the University of Bath Psychology Ethics Committee (16-148).

All volunteers working for the project were sent an information sheet inviting them to participate in an interview about their experience of working with hoarders through the hoarding project. Clients of the project were approached by the volunteer currently working with them or sent a letter inviting them to participate in an interview by post. All participants consented to take part in the interview and completed a demographic questionnaire.

To establish clutter severity, volunteers were also asked to complete the Clutter Image Rating Scale (CIRS; Frost, Steketee, Tolin, & Renaud, 2008), a self-report measure of level of clutter in the home. The CIRS comprises nine photos of increasing clutter representing a kitchen, a living room and a bedroom. Each room is scored (1-9) and a composite score is calculated. A score of four and above is indicative of problematic clutter. To assess the presence and severity of acquisition, discarding and clutter behaviours, all clients completed the Savings Inventory Revised (SI-R; Frost, Steketee, & Kyrios, 2001), a self-report measure of hoarding severity. The SI-R is a 23 item questionnaire with a total available score of 92, a score of 40 or above is indicative of clinically severe hoarding difficulties.

### **Design**

Qualitative methods were adopted to explore participants' experiences in depth and identify new concepts (Pope & Mays, 1995). Interpretive Phenomenological Analysis (IPA; Smith & Osborn, 2008; Smith, 1996) was considered a suitable approach for this study as it captures an individual's personal experiences and recognises our experiences influence how we view the world around us (Yardley, 2000). IPA is concerned with an individual's subjective report, rather than formulating an objective account. It is a dynamic process

whereby the researcher adopts an active role to take an insider's perspective and use their own conceptions to interpret information (Smith & Osborn, 2008; Smith, 1996).

### **Data collection**

All interviews were audio recorded with participants' consent and lasted between 40 to 60 minutes. The interviews were transcribed verbatim and anonymised (V1-7; C1-4) to ensure confidentiality and privacy of participants. The interviews were semi-structured and aimed to elicit information about volunteers' experiences of providing and clients' experiences of receiving help. An interview schedule with open-ended questions was developed prior to the interviews using relevant research literature and was used to facilitate the participants' ability to tell their story in their own words, a central premise of IPA. Participants were encouraged to speak freely and openly about their experience and therefore played a central role in the interview.

### **Analysis**

The data from the qualitative interviews was manually analysed using IPA and classified and coded by key concepts, themes and emerging categories (Smith & Osborn, 2008). The background and stance of the research team was cognitive-behavioural, and interpretations of the interview data were taken from this position. Each interview transcript was repeatedly read and listened to. Initial notes and possible codes, summarising the experience described by participants, were noted in the margin. The transcripts were re-read and theme titles that emerged were noted. To maintain the phenomenological nature of IPA, key phrases and words from the interview data were used to describe themes. The same process was followed to analyse each interview and new theme clusters were checked against the original transcripts as they emerged to ensure foundation in participants' narratives and shared understanding. Themes were updated by shared meaning as the analysis progressed and a number of superordinate themes were identified.

Themes with low frequency across participants or weak evidential base were discarded. One volunteer talked about her difficulty maintaining boundaries without insulting her client or damaging the therapeutic relationship. She described how her client was very generous and would often make attempts to give her gifts. However, this contradicted the charity's professional boundaries as volunteers are unable to accept gifts. Another volunteer described feeling disappointed and hopeless following an unsatisfactory ending with her client when their work together ended abruptly. Although these themes had low frequency across participants and were therefore discarded, they provide an insight into the impact of the work on volunteers.

A number of validation methods and credibility checks were adopted. Regular meetings were held by the research team to ensure transparency of process and collectively move from data collection to final interpretations. The research team individually reviewed a sample of the transcripts which further helped the primary researcher to reflect on the development of themes. In keeping with the principles of IPA, the analysis presented is the researchers' interpretation of the data and other interpretations and perspectives are entirely possible.

## Results

### Sample Characteristics

All participants identified as White UK. The seven volunteers' ages ranged from 32-84 years and the four clients' ages ranged from 63-88 years. All clients identified that they had a problem accumulating and discarding numerous possessions and had never sought professional help for their difficulties. Volunteers had worked in the project for an average of eleven months (ranging from 4-24 months) and had worked with an average of two clients each (ranging from 1-4). Only one volunteer had previous experience of working with hoarders, describing their experience as "limited", and all volunteers had completed a half-day hoarding training as part of their project induction. Additional sample characteristics are provided in Table 2.1.

Table 2.1. *Sample Characteristics of clients and volunteers (N = 11)*

	<b>Volunteers</b>	<b>Clients</b>
	Mean (SD) or N (%)	Mean (SD) or N (%)
<b>Age (years)</b>	52.6 (21.4)	75.8 (12.6)
<b>Gender</b>		
Male	2 (28.6 %)	2 (50.0 %)
Female	5 (71.4 %)	2 (50.0 %)
<b>Employment Status</b>		
Retired	4 (57.1 %)	4 (100.0 %)
Full-time	2 (28.6 %)	-
Homemaker	1 (14.3 %)	-
<b>Marital Status</b>		
Married	-	2 (50.0 %)
Divorced	-	1 (25.0 %)
Widowed	-	1 (25.0 %)
<b>Education</b>		
GCSE or equivalent	-	1 (25.0 %)
A-Level or equivalent	2 (28.6 %)	-
Bachelor's degree	3 (42.9 %)	-
Master's degree	2 (28.6 %)	2 (50.0 %)
Other	-	1 (25.0 %)



Ratings on the CIRS (Frost et al., 2008) confirmed that clients who volunteers had worked with and referred to were compulsive hoarders. A score of four or more on the CIRS was reported by volunteers across the three room areas of kitchen, living room and bedroom (ranging from 4-9). Three of the four clients interviewed scored in the clinically significant range on the SI-R (Frost et al., 2001) at the time of interview; however, it is noted that clients reported their scores would have been higher at the start of their contact with volunteers.

### **Qualitative Analysis**

Four superordinate themes were developed: relationship between client and volunteer; ‘live life again’; challenges; and supporting volunteers (see Table 2.2). All eleven participants provided example quotations of themes, ensuring adequate coverage across participants. Each theme will be discussed in turn using direct quotations to illustrate them.

Table 2.2. *Superordinate and Subordinate themes*

SUPERORDINATE THEME	SUBORDINATE THEME
Relationship between client and volunteer	Space to talk Non-professional status Client led
Live life again	Making space Company and quality of life Domino effect
Challenges	Shame and embarrassment Discarding possessions Uncertainty
Supporting volunteers	Training needs Peer support

#### ***Relationship between client and volunteer***

Volunteers described that “*a lot of the early visits were in effect building up a relationship*” (V1), as well as a consensus that helping the client “*wouldn’t happen if the relationship wasn’t there*” (V7). Clients and volunteers both highlighted the importance of there being space to talk, where clients had someone to talk to and felt heard:

*We spent a lot of time just, talking really, not necessarily about her belongings, but just talking in general. And, I think that, that helped her...I just listened...we had a bit of a laugh together. (V5)*

*Having someone to talk to certainly makes you feel an awful lot better about going through everything and facing one's demons or one's embarrassment. (C1)*

*We sit and chat...I find (volunteer) coming to chat just as important, just as helpful as the work (volunteer) does, because I can talk...we get on well, it's very easy, I can be myself with (volunteer). (C2)*

Shared interests or experiences, as well as appropriate self-disclosure and an open and personable style from volunteers helped to build a relationship with clients:

*Having that relationship and similar life experiences, we've been able to engage with each other...I do think that me being myself helps a lot. (V7)*

*Our interests are very similar. I love doing crafts as well so when she is talking about making things it is something that we can share, and I think all those things are important to her. (V2)*

Volunteers and clients emphasised the importance of their informal and 'non-professional' status in their ability to build a relationship and trust with clients:

*I think being volunteers...I think they kind of appreciate that we're there because we want to be, not that it's our job...it's that kind of approach we take where it is really kind of informal, it's not official, there's not a huge gap between us and them so that helps with the trust. (V7)*

*There's a separation between us and the local authority, or health professionals, mental health professionals, because they sometimes see those people as the boss or the people who are putting pressure on, the people in charge. They see us as more of a friendly face. (V3)*

*I would say I trust (the charity). Big word, you know, trust, you can't apply it to many things. (C4)*

The ability of volunteers to be flexible and take their time was also identified as an important part of the relationship that developed:

*I think the fact that we use volunteers. We're not on a time schedule...I think that is the joy of volunteers! We have the time or we can make the time whereas somebody in paid employment, there is only so much they can do. (V7)*

*We are not a professional...whether you are a doctor or a psychiatrist or psychologist you come with a label, so we're not a professional...and I think we just have time and flexibility, which sadly professionals don't because they have so many other things to do. (V5)*

A further aspect of the relationship between client and volunteer was clients feeling able to take the lead and be more in control of the therapeutic process:

*We sit down, have a chat and I'll then say "now come on (volunteer), we'll do something"...And then (volunteer) comes and does a bit more with me. Always with me, always with...(volunteer) never pushes me. (C2)*

*I accepted that I was a hoarder...I was up for it...It's always me to say "enough of the chat, let's get on with something"...so it's very much led by me and it's never, it's never led by what (volunteer) thinks we ought to do or what I should give away...it's all written down on forms as to where I want everything to go so I'm in control, which is brilliant. (C1)*

This stance was echoed by volunteers, who identified that "you are very much guided by the client" (V6) and "we are there specifically to help them at their own pace...we haven't got targets to meet". (V7)

### ***'Live life again'***

Participants described how the project "enables people to live their lives again...to get to a better place in their lives...to have hobbies...and make friendships" (V3). One client described how the support he received "means we can stay in our home" (C2). Others described how the support from volunteers enabled clients to make space:

*The most helpful thing has been being able to enjoy her home again...it's changed her life really I think...it was having huge impacts on her mental health and her mental wellbeing and being able to walk around and kind of have that clear space has given her that clear mind...she doesn't feel the need now to buy things and collect things and fill that space again. (V3)*

*I mean all of these bags and stuff have been sorted through and it's all here so I can get to things...It's a way of sorting out life...it's sorting out your life as well as your*

*belongings...I'm aware that it's going to be a long process, I mean, there is a lot of other stuff that is going to have to go and be sorted through. But there is now light at the end of the tunnel, because the journey has been started. (C1)*

In addition to making space, clients described feeling supported and noted improvements to their overall psychological wellbeing and quality of life as a result of input from volunteers:

*The difference is amazing...they have made an incredible psychological difference...it has given me back my, some self-respect. It's made me more positive. It's given me a purpose in life, it has potentially given me a way of making money...I can now stand up for myself more, it has given me confidence, self-confidence as well as self-respect...it's been life changing. It has been incredible. (C1)*

*It means sanity really...in terms of me, the impact on me is massive...you would go under, you would go under somehow (C4)*

*It helps us to cope with work I couldn't do on my own. If I tried to do this, well, I'd never get through! (C2)*

Participants described how support from volunteers created a domino effect, enabling clients to address problems in other areas of their lives:

*It has had such a wonderful knock on effect because she's being really resilient and is sorting out a lot of different problems in her life. (V2)*

*It's been a learning curve for me as well...I'm learning that, things that used to worry me, even a few months ago, I find I can just take it in my stride, keep things in perspective. (C2)*

*It enabled her to start something that she actually wanted to do but didn't know how to do it. (V3)*

*It's made a massive difference...I have gone through a stage of evolution really, so originally it was the decluttering...I've had to sort of move another area in my mind...instead of just decluttering, during the period you have been helping me, I haven't gone under. (C4)*

Volunteers also described a sense of fulfilment and achievement from their work supporting clients, as well as a motivation to address difficulties in their own lives:

*I am a people person, so, you know, I do, I do get quite a lot out of it...I do find it satisfying, rewarding...and exciting. (V5)*

*It makes me happy because I'm retired and I like something to do...we do it because we feel we are doing something useful and, it keeps us busy...I like to be involved and doing things so, that ticks that box rather well. (V1)*

*I enjoy it...it's exciting and I actually am interested in, sort of, the Psychology side of it...It has motivated me to do more clearing, so yes, I'm probably, yeah I'm gradually reducing the amount of stuff that I have, which is good. (V6)*

### **Challenges**

Participants described feelings of shame and embarrassment about the state that their home was in, as well as experiencing challenges with the physical environment that hindered the development of the therapeutic relationship:

*I was so, ashamed and embarrassed about the state that it was in. And I used to worry for days ahead about people, who were coming in...I would start panicking seriously up to a week ahead, and hardly sleep the night before. (C1)*

*When I actually came to work with a client, I probably hadn't appreciated some of the, it's not health, but sort of hygiene aspects to the work...the moth infestations, it was a real challenge at times. (V6)*

All participants reported difficulties in discarding possessions, including difficulties with access and having enough room to work:

*Getting in the house! To manoeuvre into the front door to go into the room, the access to the room was difficult. (V4)*

*I had a bag for recycling, a bag for charity, which actually from a physical point of view is very difficult when there isn't much space. So to find enough room, to put the bags, to collect all the stuff in, and actually to just take the first suitcase out... (V6)*

*In terms of me clearing stuff out, it's sporadic in a way, it's quite difficult for me to concentrate on certain areas because it is everywhere really. (C4)*

Letting go of possessions was also challenging for clients and was exacerbated by the physical limitations of some clients and volunteers:

*He couldn't actually manoeuvre things...There were practical difficulties because of his stroke...my hearing isn't absolutely perfect so it was slightly difficult to engage with him. (V1)*

*Her main barrier was her health and her ability to move, so trying to encourage her to make space during the week when I wasn't there is not something that she could do. So I think her health and mobility was, yeah, a real barrier for her. (V7)*

*The thing that makes it difficult is actually getting rid of belongings...it's so traumatic...it is just so difficult for them, they just get so panicky and anxious...it was just too overwhelming for her...I think the challenging thing was, that she did find it very painful to let go of stuff. (V5)*

However, one client described how her difficulties lay in finding a home for her discarded belongings:

*I had a pile of VHS tapes and you can't get rid of them. They charge you, the shops won't touch them. (C3)*

Finally, volunteers provided a number of examples of uncertainty about their work with clients and feelings of being kept 'at arm's length':

*I never really knew whether I was going to get in or not...never quite sure if I would be let in...I was never really that confident, you know, of carrying on...The other frustrating thing is that she would move stuff around, so I was never quite sure what was happening! (V5)*

*What I found challenging was whether what they were saying was true. (V1)*

*I sort of found it a little, sort of, frustrating that she did in once sense keep me at arm's length...she was very suspicious about things. (V5)*

*Her paranoia about certain things...It's taken a long time to get that trust with her. And that was a barrier but that's something I had to just kind of keep plugging away at. (V2)*

### ***Supporting volunteers***

Volunteers described a desire for additional training on the psychological aspects of working with compulsive hoarders, as well as training on how to manage some of the more practical aspects of the work:

*Having had practical experience, there are aspects that would be useful to include like manual handling, talking about sort of the hygiene aspects...I think probably we could do with a little more awareness or training in terms of how we deal with it.* (V6)

*I'd like a little bit of training around the psychology of everything I think... just some guidance around those things, about how to deal with someone in this situation.* (V3)

They also described the possibility of more joint working, including the possibility of developing a peer support group amongst volunteers:

*I think that's quite important, two volunteers to go together. It would help because um, they may see things you don't...if you've got somebody who can go if you can't go for instance...two people see things differently.* (V4)

*I think that is absolutely key to the volunteers, to be able to offload it...There's a lot of Psychology involved in all this. It would be wonderful if there was opportunity for that sort of discussion.* (V1)

## **Discussion**

Little attention has been given to the experiences of individuals with hoarding disorder receiving help and those that provide help. Understanding the factors that act as barriers and facilitators in supporting people with hoarding disorder may be an important part of the solution to supporting people and increase service provision in the future. To this end, four key themes were identified in the current study. The relationship that formed between client and volunteer was crucial in providing a trusting foundation from which clients felt able to move forward. Volunteers provided a space for clients to talk, and reported that appropriate self-disclosure helped to build a relationship. The informal and 'non-professional' status of volunteers enabled clients to take the lead and feel more in control of the therapeutic process. Volunteer flexibility and lack of time constraints were also seen as important and contributed to clients 'making space' in their lives and in their homes. The support from the volunteers enabled clients to 'live life again' and created a domino effect, bringing about improvements in other areas of both of their lives. Shame and embarrassment represented a significant barrier to clients, as well as practical challenges in doing the work through lack of space or physical limitations. The need for providing support for the volunteers was also identified.

The current findings mirror previous research highlighting the importance of the therapeutic relationship (Ayers et al., 2012). Having space to talk, where clients had someone to talk to and felt heard was helpful in clients feeling less isolated. Kim, Steketee, and Frost (2001) reported that hoarders often experience social isolation and therefore input from projects such as the one described in the present study may help to broaden clients' horizons and reduce their sense of isolation. Switching the focus from hoarding and associated behaviours to the client as a person and their interests may also create a space for change and bring about improvements in clients' quality of life.

Although the importance of developing a therapeutic relationship is not new, this study highlights the significance of the 'non-professional status' of volunteers, which is consistent with previous findings that non-professionals may be helpful in enhancing clients' physical and mental wellbeing (Taft et al., 2011; Uhm et al., 2016). This has significant implications for services where their very status as 'professional' may act as a barrier to people with hoarding disorder engaging with them. Statutory services could therefore look to collaborate with third sector/charity organisations to offer peer/volunteer support alongside mainstream services. Evidence from the current research highlights benefits to both clients and volunteers by working in this way. However, the study highlighted the importance of ensuring volunteers feel supported in their work and mentoring or peer support may help to facilitate this. Moreover, the findings indicate the need for all people providing hoarding support to receive comprehensive training relevant to their role and highlights the importance of continued involvement of competent others to support and supervise the work of volunteers.

Control is an important factor in hoarding disorder (Raines, Oglesby, Unruh, Capron, & Schmidt, 2014) and the findings of this study provide further support. Clients feeling in control enabled them to feel able to start sorting and discarding their possessions. Volunteers in this study were also sensitive to the need of their clients to take the lead and feel in control regarding their possessions. This may help to both understand the reluctance amongst hoarders to seek help from formal services (Pertusa et al., 2010) and the importance of the 'volunteer' status in enabling a helping relationship to develop. Compared to working with mental health, social care, housing and environmental health services, for example, volunteers are less likely to be perceived as being in a position to threaten or remove the control that people with hoarding disorder have over their possessions. This is a difficult problem for services to solve. However, the current research highlights the importance of taking time to build a therapeutic relationship and that appropriate self-disclosure and discussion regarding shared interests may help.



Professionals may therefore benefit from making changes to their practice to adopt this approach; however, these changes are contingent on service providers recognising the value in this approach and creating space for professionals to work in this way.

It is understandable given the complexity and challenges associated with compulsive hoarding that many professionals struggle in their work (Holden et al., 2016). Although compulsive hoarders are widely considered to be a hard to reach and complex group, this study revealed positive outcomes when using volunteers in the absence of recognised cognitive behavioural strategies that have previously been highlighted as important (Ayers et al., 2012). A minimum of 50% reduction in CIRS scores was found in this study from first contact with clients to when volunteers were interviewed. Although this finding is not conclusive in itself, it suggests that non-professionals with a little training can produce positive outcomes with conventional outcome measures. Finally, as clients in this study were aged 63-88 years old, the findings have implications for professionals, services and organisations who work with older adults. Eckfield and Wallhagen (2013) reported significantly higher rates of hoarding in adults over 55 years old and according to service providers, hoarding behaviour posed a serious physical threat in 81% of their elderly clients (Kim, Steketee & Frost, 2001). The findings of this study highlight that taking time to build a therapeutic relationship, where older adults are able to take the lead and feel more in control of the therapeutic process, may help to reduce hoarding behaviours and lead to more positive outcomes in the longer term.

The present study has a number of methodological limitations. There was a dominance of female volunteers in the sample but this reflected the demographic of the volunteer pool. In light of the small sample (eleven participants) and recruitment from one organisation, the findings may have limited generalisability. Purposive sampling was adopted as the study aimed to explore the experiences of providing and receiving help within a small UK-based charity; however, it may be argued that those who agreed to participate were more motivated to share their experience and clients experiencing greater shame and embarrassment may have unintentionally been excluded (Kellett, 2007). In spite of adopting a number of validation methods and credibility checks, the analysis presented is the researchers' interpretation of the data and other interpretations and perspectives are possible. A strength of the study is that all participants completed established hoarding measures of either the SI-R (Frost et al., 2001) or CIRS (Frost et al., 2008) to confirm the presence of compulsive hoarding. This study is also the first study to directly explore the reciprocal experience of compulsive hoarders receiving and volunteers providing help. It is

a further strength that this study demonstrated an alignment in experience between the four clients interviewed and the volunteers who were working with them.

### **Conclusions and Implications**

Compulsive hoarding continues to present significant challenges (Tolin et al., 2012), as well as economic, familial and social burden (Frost et al., 2000). The present study offers insights into how this could be improved and in particular highlights the importance of taking time to develop a therapeutic relationship and adopting a flexible and informal approach. By adopting this approach, this study found that volunteers with minimal training and people with hoarding disorder can develop meaningful and supportive relationships.

## References

- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders: DSM-5* (5th ed.). Washington, DC.
- Ayers, C. R., Bratiotis, C., Saxena, S., & Wetherell, J. L. (2012). Therapist and patient perspectives on cognitive-behavioral therapy for older adults with hoarding disorder: A collective case study. *Aging & mental health, 16*(7), 915-921.
- Ayers, C. R., Saxena, S., Golshan, S., & Wetherell, J. L. (2010). Age at onset and clinical features of late life compulsive hoarding. *International journal of geriatric psychiatry, 25*(2), 142-149.
- Ayers, C. R., Wetherell, J. L., Golshan, S., & Saxena, S. (2011). Cognitive-behavioral therapy for geriatric compulsive hoarding. *Behaviour research and therapy, 49*(10), 689-694.
- Chartered Institute of Environmental Health. (2009). Professional practice note: Hoarding and how to approach it – guidance for environmental health officers and others. Retrieved from [https://www.cieh.org/uploadedfiles/core/policy/publications\\_and\\_information\\_services/policy\\_publications/publications/hoarding\\_ppn\\_may09.pdf](https://www.cieh.org/uploadedfiles/core/policy/publications_and_information_services/policy_publications/publications/hoarding_ppn_may09.pdf)
- Coles, M. E., Frost, R. O., Heimberg, R. G., & Steketee, G. (2003). Hoarding behaviors in a large college sample. *Behaviour research and therapy, 41*(2), 179-194.
- Damecour, C. L., & Charron, M. (1998). Hoarding: a symptom, not a syndrome. *The Journal of clinical psychiatry, 59*(5), 267-272; quiz 273.
- Eckfield, M. B., & Wallhagen, M. I. (2013). The synergistic effect of growing older with hoarding behaviors. *Clinical nursing research, 22*(4), 475-491.
- Frost, R., Steketee, G., & Kyrios, M. (2001). Assessing the severity of compulsive hoarding: The Saving Inventory-Revised. *Unpublished manuscript*.
- Frost, R. O., & Gross, R. C. (1993). The hoarding of possessions. *Behaviour research and therapy, 31*(4), 367-381.
- Frost, R. O., Steketee, G., Tolin, D. F., & Renaud, S. (2008). Development and validation of the clutter image rating. *Journal of Psychopathology and Behavioral Assessment, 30*(3), 193-203.

- Frost, R. O., Steketee, G., & Williams, L. (2000). Hoarding: a community health problem. *Health & social care in the community*, 8(4), 229-234.
- Frost, R. O., Tolin, D. F., & Maltby, N. (2010). Insight-related challenges in the treatment of hoarding. *Cognitive and Behavioral Practice*, 17(4), 404-413.
- Grisham, J. R., Frost, R. O., Steketee, G., Kim, H.-J., & Hood, S. (2006). Age of onset of compulsive hoarding. *Journal of Anxiety Disorders*, 20(5), 675-686.
- Holden, K., Kellett, S., Davies, J., & Scott, S. (2016). The experience of working with people that hoard: a Q-sort exploration. *Journal of Mental Health*, 1-7.
- Kellett, S. (2007). Compulsive hoarding: a site-security model and associated psychological treatment strategies. *Clinical Psychology & Psychotherapy*, 14(6), 413-427.
- Kim, H.-J., Steketee, G., & Frost, R. O. (2001). Hoarding by elderly people. *Health & Social Work*, 26(3), 176-184.
- Mataix-Cols, D., Frost, R. O., Pertusa, A., Clark, L. A., Saxena, S., Leckman, J. F., . . . Wilhelm, S. (2010). Hoarding disorder: a new diagnosis for DSM-V? *Depression and anxiety*, 27(6), 556-572.
- Pertusa, A., Frost, R. O., Fullana, M. A., Samuels, J., Steketee, G., Tolin, D., . . . Mataix-Cols, D. (2010). Refining the diagnostic boundaries of compulsive hoarding: a critical review. *Clinical Psychology Review*, 30(4), 371-386.
- Pope, C., & Mays, N. (1995). Qualitative research: reaching the parts other methods cannot reach: an introduction to qualitative methods in health and health services research. *Bmj*, 311(6996), 42-45.
- Raines, A. M., Oglesby, M. E., Unruh, A. S., Capron, D. W., & Schmidt, N. B. (2014). Perceived control: A general psychological vulnerability factor for hoarding. *Personality and Individual Differences*, 56, 175-179.
- Slatter, M. (2007). Treasures, trash and tenure: hoarding and housing risk. *People, Place & Policy Online*, 2(1), 28-36.
- Smith, J., & Osborn, M. (2008). Interpretative phenomenological analysis. JA Smith (Ed.), *Qualitative psychology: A practical guide to research methods* (51–80): Los Angeles, CA: Sage.

- Smith, J. A. (1996). Beyond the divide between cognition and discourse: Using interpretative phenomenological analysis in health psychology. *Psychology and health, 11*(2), 261-271.
- Stekete, G., & Frost, R. (2007). *Compulsive Hoarding and Acquiring. Therapist Guide*. New York: Oxford University Press.
- Taft, A. J., Small, R., Hegarty, K. L., Watson, L. F., Gold, L., & Lumley, J. A. (2011). Mothers' AdvocateS In the Community (MOSAIC)-non-professional mentor support to reduce intimate partner violence and depression in mothers: a cluster randomised trial in primary care. *BMC public health, 11*(1), 178.
- Tolin, D. F., Frost, R. O., & Steketee, G. (2012). Working with hoarding vs. non-hoarding clients: A survey of professionals' attitudes and experiences. *Journal of Obsessive-Compulsive and Related Disorders, 1*(1), 48-53.
- Turner, K., Steketee, G., & Nauth, L. (2010). Treating elders with compulsive hoarding: a pilot program. *Cognitive and Behavioral Practice, 17*(4), 449-457.
- Uhm, S. Y., Tsoh, J. Y., Mackin, R. S., Gause, M., Chan, J., Franklin, J., . . . Bain, D. (2016). Comparison of a peer facilitated support group to cognitive behavior therapy: Study protocol for a randomized controlled trial for hoarding disorder. *Contemporary clinical trials, 50*, 98-105.
- Yardley, L. (2000). Dilemmas in qualitative health research. *Psychology and health, 15*(2), 215-228.

Main Research Project

**Posttraumatic growth in mothers 2-6 years after stillbirth and early miscarriage**

Kirsty Ryninks

Doctorate in Clinical Psychology

Department of Psychology, University of Bath, Claverton Down, Bath, BA2 7AY

Email: K.E.Ryninks@bath.ac.uk

Word count: 5318

May 2018

**Internal/academic supervisors:** Dr Megan Wilkinson-Tough; Doctoral Programme  
in Clinical Psychology

Department of Psychology, University of Bath, Claverton Down, Bath, BA2 7AY

Email: M.J.Wilkinson-Tough@bath.ac.uk

**External/field supervisors:** Professor Antje Horsch, Clinical Psychologist; Institute  
of Higher Education and Research in Healthcare, University of Lausanne and  
Department Woman-Mother-Child, University Hospital Lausanne, Switzerland

Email: Antje.Horsch@chuv.ch

Dr Sarah Stacey, Clinical Psychologist; HMP Bristol, 19 Cambridge Road, Bristol,  
BS7 8PS

Email: sarah.stacey1@nhs.net

**Proposed Journal:** BMC Pregnancy and Childbirth

Please see Appendix VII for instructions to authors



## Introduction

Losing an unborn child is a very painful experience for a parent and may cause more intense grief than the loss of a partner or parent (Bonanno et al., 2002). Mothers may experience a variety of losses in pregnancy, including early miscarriage (EM; occurring in the first trimester of pregnancy up to 12 weeks), late miscarriage (LM; occurring during the second trimester at 13-23 weeks) and stillbirth (SB; defined as intrauterine death after 24 weeks gestation in the United Kingdom (UK)). Miscarriage affects 200,000 couples every year in the UK and 85% of all miscarriages occur in the first trimester. The global prevalence rate of SB in 2015 was 2.7 million (Lawn et al., 2016) and SB is estimated to occur in nearly 1 in 200 pregnancies (Martin, Kochanek, Strobino, Guyer, & MacDorman, 2005). In comparison to other types of child loss, pregnancy loss is often treated as less significant (Frøen et al., 2011) and parents are left feeling isolated and invalidated in their grief.

Losing a baby through miscarriage (Lee & Slade, 1996) or SB (Horsch, Jacobs, & McKenzie-McHarg, 2015) is recognised as traumatic (Kersting & Wagner, 2012). Part of the future is lost when a baby dies and bereavement reactions have been reported to be pervasive, powerful and complex (Cacciatore, 2010; Campbell-Jackson & Horsch, 2014; Robinson, 2011). Mothers are often unprepared for the loss and experience a range of psychological reactions, including denial, guilt, anger, grief and feelings of ‘empty arms’ (Adeyemi et al., 2008; Brownlee & Oikonen, 2004; Cacciatore, 2010; Murphy, Shevlin, & Elklit, 2014). Although the majority of mothers adjust and regain a sense of purpose, 15-25% seek additional support for enduring adjustment difficulties in the year after their loss (Bennett, Litz, Lee, & Maguen, 2005; Conway & Russell, 2000). In comparison to mothers of live babies, higher depressive and anxiety symptoms are more likely to be experienced by mothers for up to three years following the loss of their baby (Cacciatore, Rådestad, & Frederik Frøen, 2008). Mothers are also at risk of developing post-traumatic stress (Engelhard, van den Hout, & Arntz, 2001; Horsch et al., 2015), and experiencing high levels of affective disturbance in a subsequent pregnancy (Turton, Hughes, Evans, & Fainman, 2001).

Most studies to date have grouped different perinatal losses in their efforts to understand the psychological impact of loss in pregnancy. However, there are a number of differences between EM and SB and studies have highlighted that the psychological impact of these differing losses requires further clarification (Cacciatore & Bushfield, 2007; Klier, Geller, & Ritsher, 2002). Evidence suggests the further a pregnancy has progressed, the greater the attachment to the unborn child and more severe the grief reaction experienced by



the mother (Goldbach, Dunn, Toedter, & Lasker, 1991). Where loss in pregnancy is experienced early on, the loss often remains unacknowledged outside of the immediate family (Cacciatore, 2013; Kersting & Wagner, 2012). In contrast, as SB occurs after 24 weeks gestation, the pregnancy is visible due to significant physical changes in the appearance of the mother. In order to develop a deeper understanding of the psychological impact of miscarriage and SB, it is important to compare both losses within the same study (Cuisinier, Kuijpers, Hoogduin, De Graauw, & Janssen, 1993) and grouping perinatal loss is unlikely to be helpful (Campbell-Jackson & Horsch, 2014). Although this has been done by a few studies previously, findings are not consistent. For example, Toedter, Lasker, and Alhadeff (1988) found grief was more intense with greater gestational age at loss, whereas Smith and Borgers (1989) found no differences in grief between women who had experienced miscarriage compared to SB or neonatal death.

Apart from the distressing emotions that follow traumatic events, individuals may also experience positive psychological change (Tedeschi & Calhoun, 2004b). Posttraumatic Growth (PTG) is defined as “positive psychological change experienced as a result of the struggle with highly challenging life circumstances” (Calhoun & Tedeschi, 2001, p. 157). Through the journey to make sense of the world post-trauma, individuals may develop a greater appreciation for their relationships, have new spiritual insights, recognise their strengths and develop an acceptance that it is not always possible to prevent negative outcomes. Posttraumatic growth has “a quality of transformation or qualitative change in functioning” (Tedeschi & Calhoun, 2004b) and goes beyond an individual’s previous adaption, psychological functioning or life awareness. Any positive changes arising in response to the trauma occur in conjunction with grief and distressing emotions that follow traumatic events, not in place of them (Tedeschi & Calhoun, 2004b). The phenomenon of PTG has been reported in individuals who have faced a variety of traumatic circumstances, including: natural disasters (Cryder, Kilmer, Tedeschi, & Calhoun, 2006), terrorism (Laufer & Solomon, 2006), cancer (Bellizzi, 2004), childhood sexual abuse (Wright, Crawford, & Sebastian, 2007), burn injuries (Rosenbach & Renneberg, 2008), war (Maugen, Vogt, King, King, & Litz, 2006), road trauma (Harms & Talbot, 2007) and HIV/AIDS (Milam, 2006). While the death of a loved one is devastating and painful, coping with bereavement has also been found to provide a context for significant positive change (Calhoun & Tedeschi, 1990; Engelkemeyer & Marwit, 2008; Mathews & Servaty-Seib, 2007). The possibility of PTG in the aftermath of pregnancy loss has received limited attention to date, with a focus historically on the more traumatic and distressing aspects of this loss. Black and Wright (2012) highlighted the need for systematic research on different perinatal losses (e.g., EM and SB) to clarify the posttraumatic responses these losses engender in bereaved parents.

One of the first studies to investigate PTG after pregnancy loss was conducted by Büchi et al. (2007). They investigated PTG in 54 parents who had lost their premature baby within the previous two to six years. Parents reported a shift in priorities about what is important in life, with the change appearing greater in mothers (78%) than fathers (44%). Wright (2010) used grounded theory with a sample of nineteen women who had experienced loss in pregnancy and found women felt more loving, compassionate and appreciative of the relationships they had. Black and Sandelowski (2010) conducted an ethnographic study to explore personal growth following severe fetal diagnosis and described that women and their partners found a new appreciation for life following their loss. More recently, Krosch and Shakespeare-Finch (2017) reported moderate levels of PTG ( $M=51.22$ ,  $SD=20.13$ ) on the Posttraumatic Growth Inventory (Tedeschi & Calhoun, 1996) in women who had experienced pregnancy loss. However, so far, PTG has not been investigated in detail following pregnancy loss and no study has explicitly compared PTG in SB and EM.

Calhoun, Tedeschi, Cann, and Hanks (2010) presented a Model of Growth in Grief to understand how losing “a close other” may result in the acknowledgment of a number of positive changes. Models of PTG focus on the degree to which the traumatic experience challenges world assumptions (Janoff-Bulman, 2006; Tedeschi & Calhoun, 2006). Deaths that are unexpected, less ‘natural’ and conflict with or disrupt an individual’s assumptions and beliefs about the way the world should work (e.g. the death of a child) often result in more distress and PTG (Tedeschi & Calhoun, 2006). In addition to dealing with distress from losing a close other, the individual must reassess the principles of how they believe the world should work (Calhoun et al., 2010). Rumination has an important role to play as individuals attempt to reconstruct their beliefs following bereavement. Whilst this process can lay a foundation for growth and process of rebuilding (deliberate rumination), it can also initially add to distress (intrusive rumination) and negatively impact PTG (Calhoun & Tedeschi, 2006). Disclosure plays an important role, with supported self-disclosure helping individuals to manage distress and rebuild challenged assumptions about death and how the world should work (Calhoun et al., 2010).

The Model of Growth in Grief is of particular interest when considering loss in pregnancy, as the majority of published findings of positive changes following loss have focused on the loss of a loved one with whom one had an ongoing and longstanding relationship. Although mothers form attachments early on in pregnancy (Gold, Sen, & Hayward, 2010), mothers who experience loss in pregnancy do not have the opportunity to create direct life experiences with their baby, unlike in the death of close family members

(Kersting & Wagner, 2012). The pathway to PTG may therefore differ to other types of loss. Moreover, although mothers following EM and SB both experience loss in pregnancy, particular factors relating to the Model of Growth in Grief (challenge to assumptive beliefs (CAB), disclosure) may differ and result in differing levels of PTG. To our knowledge, no study has so far tested the Model of Growth in Grief using between-group comparisons. This study aimed to investigate PTG in mothers two to six years following SB compared to EM. It was hypothesised that mothers following SB will demonstrate more (1) PTG, (2) CAB, and (3) disclosure than mothers following EM. The study also aimed to investigate how theoretically-derived variables of the Model of Growth in Grief (CAB, rumination and disclosure) explain unique variance in PTG when key factors are controlled.

## **Method**

### **Design**

A cross-sectional between-groups design was adopted to compare PTG in mothers following SB and EM. The independent variable was type of loss (SB/EM) and primary dependent variable was PTG, with secondary dependent variables identified as key components from the Model of Growth in Grief (CAB, rumination and self-disclosure). The study was advertised and completed online. Ethical approval was granted by the University of Bath Psychology Ethics Committee (17-044).

### **Participants**

The sample comprised 57 mothers in the SB group and 63 mothers in the EM group. Women were eligible to participate if they spoke English, were aged over 18 years old and had experienced either a SB at 24 weeks gestation or later (UK definition; Hughes & Riches, 2003) or an EM before 13 weeks gestation in the last two to six years. This time point was used previously by Büchi et al. (2007) and was established as sufficient time for PTG to have occurred. Women who had experienced a late miscarriage (between the 13<sup>th</sup> and 23<sup>rd</sup> week of pregnancy) were not included in this study due to concerns regarding feasibility, time constraints and greater heterogeneity across late miscarriage than in the EM or SB populations. Women were recruited online through advertisements placed on social media sites and through social media accounts of perinatal death charities. Data collection took place between November 2017 and March 2018.

### **Procedure**

All questionnaires were completed online following reading of study information and provision of informed consent. For all questions referring to ‘the event’ or ‘the

incident', participants were asked to think about 'losing my baby'. If participants had experienced multiple losses in the previous two to six years, they were asked to answer the questions in relation to the loss (SB/EM) that affected them the most. Completing the survey took 30 minutes.

## **Materials**

### ***Emotional distress***

*Depressive symptoms.* The Patient Health Questionnaire (PHQ-8; Kroenke & Spitzer, 2002) is an eight-item measure of depression with total score range of 0-24. The PHQ-8 omits the final item of the PHQ-9 which relates to thoughts about death or self-harm. It has comparable characteristics to the nine item version for diagnosing depressive disorders, with good test-retest reliability ( $\alpha=.84$ ) (Kroenke & Spitzer, 2002; Kroenke, Spitzer, & Williams, 2001; Kroenke et al., 2009). Cronbach's  $\alpha$  for this study was .92.

*Anxiety symptoms.* The Generalised Anxiety seven-item scale (GAD-7; Spitzer, Kroenke, Williams, & Löwe, 2006) is a seven-item measure of anxiety with total score range of 0-21. The GAD-7 has good reliability, as well as criterion, construct, factorial and procedural validity (Spitzer et al., 2006). Cronbach's  $\alpha$  for this study was .91.

*Grief symptoms.* The Perinatal Grief Scale Short Version (PGS; Potvin, Lasker, & Toedter, 1989) is a 33-item scale with total score range of 33-165. Higher scores reflect more grief. Internal reliability of the total scale is excellent (Potvin et al., 1989), Cronbach's  $\alpha=.94$  (this sample  $\alpha=.94$ ). For the purposes of this study, only the total scale was used.

*Posttraumatic stress disorder symptoms.* The Posttraumatic Stress Disorder Checklist for DSM-5 (Weathers et al., 2013) is composed of 20 items that correspond to the 20 criteria for PTSD outlined in DSM-5. It contains four subscales corresponding to the four symptom clusters. For each item, a score of two or above is regarded as clinically relevant. The PCL-5 has demonstrated excellent internal consistency ( $\alpha=.95$ ) and strong divergent validity (Ashbaugh, Houle-Johnson, Herbert, El-Hage, & Brunet, 2016). For the purposes of this study, only the total scale was used. Cronbach's  $\alpha$  for this study was .96.

### ***Posttraumatic Responses and Disclosure***

*Posttraumatic Growth symptoms.* The Posttraumatic Growth Inventory (PTGI; Tedeschi & Calhoun, 1996) is a 21-item scale with total score range of 0-105, where higher scores indicate more growth. It was designed to measure positive existential growth following traumatic events and consists of five subscales. In this study, the phrase 'your crisis' in the instruction was changed to 'losing your baby'. The PTGI has good internal consistency and

moderate test-retest reliability (Tedeschi & Calhoun, 1996). For the purposes of this study, only the total scale was used. Cronbach's  $\alpha$  for this study was .94

*Challenge to Assumptive Beliefs.* The Core Beliefs Inventory (Cann et al., 2010) is a nine-item scale where participants rate the degree to which a recent highly stressful event led them to re-examine a number of core assumptions about themselves and their world. The total score range is 0-45. The CBI has construct validity, acceptable test-retest reliability, and very good internal consistency (Cann et al., 2010). Cronbach's  $\alpha$  for this study was .92.

*Disclosure of Trauma.* The Disclosure of Trauma Questionnaire (DTQ; Müller, Beauducel, Raschka, & Maercker, 2000) is a 34-item self-report measure comprised of three subscales: 'reluctance to talk' (reported resistance to tell others about the trauma); 'urge to talk' (individual's need to disclose the traumatic experience); and 'emotional reactions' (descriptions of affective states experienced while disclosing the trauma). For the purpose of this study, the instructions for the DTQ were expanded to include 'in relation to losing your baby'. The DTQ has good psychometric properties, with Cronbach's  $\alpha = 0.82$  to  $0.88$  and retest reliability of  $0.76$  to  $0.89$  (Müller et al., 2000). For the current study, Cronbach's  $\alpha$  reliability coefficients for the subscales were .81 (reluctance to talk), .88 (urge to talk), .87 (emotional reactions) and .89 for the total.

*Actual Self-disclosure.* In the absence of suitable formal published measures of actual self-disclosure, a set of questions was developed by the authors. Participants were asked to estimate the number of hours at three time points (first month, first year and second year after loss) they had spent talking to others (e.g. friends, family, healthcare staff or anyone else) about their feelings about losing their baby. They were also asked whether they had talked enough about their feelings about losing their baby at these time points. Actual self-disclosure was calculated by combining the total hours participants spent talking about their feelings about losing their baby in the first and second year after loss.

*Rumination.* The Event Related Rumination Inventory (Cann et al., 2011) is composed of two subscales containing ten statements relevant to intrusive rumination (IR) and ten statements relevant to deliberate rumination (DR). The total score range is 0-60. In this study, the instructions for the ERRI were changed from 'After an experience like the one you reported' to 'After a loss in pregnancy'. The ERRI has construct validity, acceptable test-retest reliability, and very good internal consistency (Cann et al., 2011). For the current study, Cronbach's  $\alpha$  reliability coefficients for the subscales were .95 (IR), .92 (DR) and .95 for the total.

## **Statistical Analysis**

A power analysis was conducted using G\*power to calculate the sample size required to determine a statistical difference in PTG between the two groups. With an alpha of 0.05 and power of 0.80, the projected sample needed in each group was 51 participants for a moderate effect. Survey data was analysed using IBM SPSS Statistics version 23. Missing data points were identified in the actual disclosure variable, where 15 participants had not been able to estimate the amount of time they had spent talking about losing their baby. As this variable was considered very susceptible to individual differences, the missing data points were not substituted. Two outliers in actual self-disclosure were identified and excluded from analyses involving this variable. Descriptive statistics were used to describe the sample and all variables were assessed for outliers and violations of normality using histograms and skewness and kurtosis statistics. Where assumptions of normality were violated, equal variances were not assumed when testing for significance. Pearson's chi-square analyses and independent-samples t-tests were conducted to examine between-group differences that might affect or confound subsequent analyses. A hierarchical stepwise regression analysis was conducted to investigate how theoretically-derived variables of the Model of Growth in Grief (CAB, rumination, disclosure) explained unique variance in PTG when key factors were controlled for. The variables were entered in four steps in line with the development of PTG through the Model of Growth in Grief: 1) Confounds - age, type of loss (SB vs EM), PTSD symptoms and perinatal grief; 2) CAB; 3) Rumination - DR and IR; 4) Disclosure - urge to talk and actual self-disclosure.

## **Results**

### **Participant Characteristics**

Living children were reported in 96.5% of the SB group (N=55) and 87.3% of the EM group (N=55; see Table 3.1 for sample characteristics). Eleven participants in the SB group (19.3%) reported being pregnant at the time of completing the study compared to six participants in the EM group (9.5%). The mean time since loss was 3.23 years (SD=1.13) following SB and 3.19 years (SD=1.09) following EM. A total of 42.1% of participants in the SB group (N=24) and 57.1% in the EM group (N=36) had previously experienced loss in pregnancy. There were no significant differences between group and ethnicity, marital status, education, employment, being pregnant at the moment or having had a previous loss in pregnancy. However, more participants in the EM group (14.3%; N=9) compared to SB (1.8%; N=1) had previously experienced a late miscarriage ( $\chi^2(1) = 6.15, p = .013$ ). A significant difference in age between the SB (M=34.74, SD=4.71) and EM groups (M=36.71, SD=5.36;  $t(118) = -2.14, p = .035$ ) was also found.

Table 3.1 *Sample Characteristics (N = 120)*

	<b>Stillbirth</b> Mean (SD) or N (%)	<b>Early Miscarriage</b> Mean (SD) or N (%)	<b>p</b>
Age (years)	34.74 (4.71)	36.71 (5.36)	.035
Ethnicity			.392
White British	53 (93%)	57 (90.5%)	
Other White background	4 (7%)	3 (4.8%)	
Multiple/Mixed ethnic group	-	2 (3.2%)	
Other	-	1 (1.6%)	
Marital Status			.410
Single, never married	-	1 (1.6%)	
Married/civil partnership or cohabiting	56 (98.2%)	59 (93.7%)	
Divorced or separated	1 (1.8%)	3 (4.8%)	
Employment Status			.511
Full-time	14 (24.6%)	18 (28.6%)	
Part-time	26 (45.6%)	24 (38.1%)	
Student	-	2 (3.2%)	
Homemaker	16 (28.1%)	16 (25.4%)	
Other	1 (1.8%)	3 (4.8%)	
Education			.485
Left school without qualifications	-	1 (1.6%)	
GCSE qualifications or equivalent	4 (7%)	5 (7.9%)	
A-Level or equivalent	11 (19.3%)	8 (12.7%)	
Degree qualification or above	38 (66.7%)	48 (76.2%)	
Prefer not to say	1 (1.8%)	-	
Other	3 (5.3%)	1 (1.6%)	
Gestation of pregnancy loss			-
1-4 weeks	-	2 (3.2%)	
5-8 weeks	-	33 (52.4%)	
9-12 weeks	-	28 (44.4%)	
24-27 weeks	10 (17.5%)	-	
28-31 weeks	7 (12.3%)	-	
32-35 weeks	6 (10.5%)	-	
36-40 weeks	23 (40.4%)	-	
Over 40 weeks	11 (19.3%)	-	
Previous Pregnancy Loss			.100
Early Miscarriage	23 (40.4%)	34 (54%)	.136
Late Miscarriage	1 (1.8%)	9 (14.3%)	.013
Stillbirth	2 (3.5%)	2 (3.2%)	.919

*SD* standard deviation; *N* number of participants; *p* p-value

## Descriptive statistics

Mean scores and standard deviations for all measures are provided in Table 3.2. Independent-samples t-tests revealed the SB group had significantly more perinatal grief ( $t(118)=3.36, p<.001$ ) and PTSD symptoms ( $t(117.97)=2.15, p=.034$ ) compared to EM. There were no significant differences in anxiety scores ( $t(117.55)=1.38, p=.170$ ) or depression scores ( $t(117.98)=.09, p=.930$ ) between the groups.

## Posttraumatic responses

Independent-samples t-tests revealed that compared to EM, the SB group had significantly greater PTG ( $t(117.53)=3.19, p=.002$ ). Significant differences were found in IR ( $t(118)=3.70, p<.001$ ) and DR ( $t(118)=2.04, p=.043$ ) for SB compared to EM. There was also a significant difference in CAB for SB compared to EM ( $t(117.78)=5.17, p<.001$ ).

Table 3.2. Means (and standard deviations) for questionnaire scores by type of loss

	Stillbirth Mean (SD)	Early Miscarriage Mean (SD)	<i>t</i>
GAD-7 (anxiety)	8.95 (5.65)	7.49 (5.877)	1.38
PHQ-8 (depression)	7.79 (6.28)	7.68 (7.043)	.09
PGS (perinatal grief)	98.05 (24.30)	81.97 (27.720)	3.36***
PCL5 (PTSD symptoms)	29.42 (19.30)	21.51 (21.034)	2.15*
DTQ (disclosure)	42.11 (16.97)	33.44 (15.719)	2.90**
DTQ – reluctance to talk	13.16 (8.66)	11.44 (8.890)	1.07
DTQ – urge to talk	13.68 (7.36)	10.00 (7.285)	2.75**
DTQ – emotional reactions	15.26 (7.70)	12.00 (6.816)	2.46*
Actual Self-disclosure (hours)	287.79 (317.78)	38.95 (69.266)	5.32***
ERRI (rumination)	35.84 (11.90)	27.52 (16.053)	3.20**
ERRI – intrusive rumination	17.88 (6.77)	12.62 (8.575)	3.70***
ERRI –deliberate rumination	17.96 (7.34)	14.90 (8.891)	2.04*
CBI (challenge to assumptive beliefs)	24.04 (9.08)	14.79 (10.491)	5.17***
PTGI (posttraumatic growth)	45.96 (23.31)	32.11 (24.195)	3.19**

*SD* standard deviation, *t* t-value, *GAD-7* Generalised Anxiety seven item scale, *PHQ-8* Patient Health Questionnaire, *PGS* Perinatal Grief Scale Short Version, *PCL-5* Posttraumatic Stress Disorder Checklist for DSM-5, *DTQ* Disclosure of Trauma Questionnaire, *ERRI* Event Related Rumination Inventory, *CBI* Core Beliefs Inventory, *PTGI* Posttraumatic Growth Inventory.

\*  $p < .05$ , \*\*  $p < .01$ , \*\*\*  $p < .001$



## **Disclosure**

Significant differences in urge to talk ( $t(116.55) = 2.75, p = .007$ ) and emotional reactions during disclosure ( $t(118) = 2.46, p = .015$ ) were found in the SB compared to EM group. There was no significant difference in reluctance to talk between the two groups ( $t(117.34) = 1.07, p = .287$ ). Compared to EM, the SB group had significantly more actual self-disclosure ( $t(50.90) = 5.32, p < .001$ ). In the first month and year after losing their baby, a greater percentage of participants felt they did not talk enough about their EM (60.3% in first month ( $N=38$ ); 55.6% in first year ( $N=35$ )) compared to participants following SB (47.4% in first month ( $N=27$ ); 49.1% in first year ( $N=28$ )). In contrast, more participants following SB (59.6%,  $N=34$ ) reported they did not talk enough in the second year after losing their baby compared to participants following EM (46%,  $N=29$ ).

### **Variables predicting PTG**

PTG was found to be significantly correlated with type of loss,  $r(120) = -.28, p = .002$ , CAB,  $r(120) = .47, p < .001$ , DR,  $r(120) = .30, p = .001$ , urge to talk,  $r(120) = .54, p < .001$ , reluctance to talk,  $r(120) = -.19, p = .038$ , and actual self-disclosure,  $r(103) = .37, p < .001$ . The following variables were not significantly correlated with PTG: PTSD symptoms, perinatal grief, depression, anxiety, IR, emotional reactions, previous loss in pregnancy, experiencing multiple losses in the last 2-6 years, having living children, being pregnant now, having a child since experiencing a pregnancy loss, age and time since pregnancy loss (see Table 3.3 for correlations of key variables).

The results of the hierarchical stepwise regression analysis are outlined in Table 3.4. Overall, 8.2% of variance in PTG was explained from step 1 (age, type of loss, PTSD symptoms and perinatal grief). CAB predicted an additional 17.6% in step 2. IR and DR predicted 5.9% in PTG in step 3 and urge to talk and actual self-disclosure predicted a further 14.7% in step 4. The final model explained 46.4% of variance in PTG,  $F(9,93) = 8.93, p < .001$ . Urge to talk ( $\beta = .39$ ), CAB ( $\beta = .38$ ) and actual self-disclosure ( $\beta = .20$ ) significantly predicted PTG. The other variables were not found to make a unique contribution.

Table 3.3. *Correlations for SB and EM groups*

Measure	PTGI	Age	Type of Loss	Time since Loss	Child since Loss	PCL-5	PGS	CBI	ERRI Int.	ERRI Delib	DTQ Urge to Talk	DTQ Reluc to Talk
PTGI												
Age	-.059											
Type of Loss	-.28***	.19*										
Time since Loss	-.11	.08	-.02									
Child since Loss	.14	.00	-.20*	.14								
PCL-5	.076	-.10	-.19*	-.07	-.23*							
PGS	.13	-.10	-.30**	-.03	-.23*	.79***						
CBI	.47***	-.05	-.43***	-.01	-.11	.46***	.56***					
ERRI Int	.16	-.07	-.32***	-.12	-.10	.76***	.68***	.37***				
ERRI Del	.30***	.05	-.19*	-.08	-.24**	.56***	.50***	.39***	.61***			
DTQ Urge	.54***	-.04	-.25**	-.12	.001	.27**	.30**	.39***	.34***	.45***		
DTQ Reluc	-.19*	-.13	-.10	.17	-.18*	.47***	.43**	.33***	.35***	.13	-.19*	
Actual Self-Disclosure	.37***	-.09	-.49***	-.05	.20*	.06	.14	.33***	.13	.12	.20*	-.12

*PTGI* Posttraumatic Growth Inventory, *PCL-5* Posttraumatic Stress Disorder Checklist for DSM-5, *PGS* Perinatal Grief Scale Short Version, *CBI* Core Beliefs Inventory, *ERRI* Event Related Rumination Inventory, *DTQ* Disclosure of Trauma Questionnaire. \*  $p < .05$ , \*\*  $p < .01$ , \*\*\*  $p < .001$

Table 3.4. *Multiple Regression Analysis: variables included in the regression model*

	<i>B</i>	<i>SE B</i>	$\beta$	<i>t</i>	<i>p</i>
Step 1					
Constant	54.20	21.24		2.55	.01
Age	-.02	.48	-.00	-.03	.97
Type of Loss	-13.06	5.08	-.27	-2.57	.01
PTSD symptoms	-.04	.19	-.03	-.21	.83
Perinatal grief	.07	.15	.07	.46	.65
Step 2					
Constant	42.45	19.35		2.19	.03
Age	-.15	.43	-.03	-.35	.72
Type of Loss	-5.10	4.88	-.10	-1.05	.30
PTSD symptoms	-.10	.17	-.08	-.59	.56
Perinatal grief	-.13	.14	-.14	-.93	.36
Challenge to assumptive beliefs	1.22	.26	.54	4.79	.00
Step 3					
Constant	38.31	19.05		2.01	.05
Age	-.32	.42	-.07	-.76	.45
Type of Loss	-3.55	4.92	-.07	-.72	.47
PTSD symptoms	-.32	.20	-.26	-1.61	.11
Perinatal grief	-.17	.14	-.18	-1.21	.23
Challenge to assumptive beliefs	1.17	.26	.51	4.61	.00
Intrusive Rumination	.31	.44	.10	.69	.49
Deliberate Rumination	.79	.33	.27	2.39	.02
Step 4					
Constant	20.78	17.61		1.18	.24
Age	-.23	.38	-.05	-.61	.54
Type of Loss	1.18	4.79	.04	.37	.71
PTSD symptoms	-.23	.18	-.19	-1.28	.20
Perinatal grief	-.15	.12	-.17	-1.25	.22
Challenge to assumptive beliefs	.86	.24	.38	3.64	.00
Intrusive Rumination	.15	.40	.051	.39	.70
Deliberate Rumination	.36	.31	.12	1.17	.24
Urge to Talk	1.28	.29	.39	4.40	.00
Actual self-disclosure	.019	.01	.20	2.25	.03

*B* unstandardised beta, *SE B* standard error for unstandardised beta,  $\beta$  standardised beta (regression coefficient), *t* t-value, *p* probability.

Note  $R^2 = .082$  for step 1;  $\Delta R^2 = .176$  for step 2 ( $p < .001$ );  $\Delta R^2 = .059$  for step 3 ( $p < .05$ );  $\Delta R^2 = .147$  for step 4 ( $p < .001$ )

## Discussion

This study compared levels of PTG in mothers two to six years following SB or EM and examined whether theoretically-derived variables of the Model of Growth in Grief explained unique variance in PTG when key factors were controlled. Results indicated mothers who had experienced a SB experienced significantly greater PTG, PTSD symptoms and perinatal grief than mothers who had experienced an EM. Greater challenge to assumptive beliefs, intrusive and deliberate rumination, greater urge to talk and consequently greater reported actual self-disclosure were also found in mothers following SB compared to EM. The final regression model explained 46.4% of variance in PTG.

Higher levels of PTSD symptoms and PTG were found following SB compared to EM. A significant difference in PTSD symptoms was found between the two groups and gestational age has been reported to predict PTSD symptomatology (Engelhard et al., 2001). The differences in PTSD symptoms between SB and EM may also have been due to the greater physical trauma and perceived risk in SB, as well as the differences in prenatal attachment. Previous studies have reported PTG in mothers following SB (Goutaudier, Nahi, Boudoukha, Séjourné, & Chabrol, 2017), miscarriage (Isguder et al., 2017) and pregnancy loss in general (Büchi et al., 2009; Büchi et al., 2007; Krosch & Shakespeare-Finch, 2017). However, this is the first study we are aware of that explicitly compared PTG in a SB and EM population. Krosch and Shakespeare-Finch (2017) found higher levels of PTG following pregnancy loss ( $M=51.22$ ,  $SD=20.13$ ) than in the present study, which may be due to their longer time since loss ( $M=4.01$  years compared to 3.23 (SB) and 3.19 (EM) in this study). There continues to be a paucity of research on the timeframe required for PTG to develop. Black and Sandelowski (2010) identified PTG in mothers within four months of perinatal loss and the present study replicated the time frame used by Büchi et al. (2007) in their study of PTG.

No significant differences between anxiety or depression were found between the groups, with mean scores in the mild range (Kroenke & Spitzer, 2002; Kroenke et al., 2009; Spitzer et al., 2006). Participants reported moderate to high levels of perinatal grief and scores were consistent with previous research (Toedter, Lasker, & Janssen, 2001) and findings that bereaved parents can experience grief for many years following their loss (Badenhorst & Hughes, 2007; Brier, 2008). Consistent with previous research (Clauss, 2009), higher grief scores were found in the SB group, further highlighting the distressing nature of this loss.

This is the first study to test the Model of Growth in Grief using a non-correlational design. Significant differences were found between the two groups, highlighting the importance of comparing types of loss within the same study. Significant group differences regarding CAB were found, suggesting SB is associated with a greater disruption of mothers' beliefs about the way the world should work than EM. These findings are consistent with theory (Calhoun et al., 2010; Tedeschi & Calhoun, 2006) and previous research highlighting the differences in CAB by gestational age (Krosch & Shakespeare-Finch, 2017).

Over half of mothers following EM felt they did not talk enough about their loss in the first month and year following their miscarriage, mirroring previous reports that EM remains shrouded in shame and silence and the difficulty mothers have in talking about their loss (Kluger-Bell, 2000). Mothers following SB disclosed greater urge to talk and reported greater actual self-disclosure than mothers following EM. Crawley, Lomax, and Ayers (2013) emphasised the importance of sharing memories of the stillborn baby to aid psychological adjustment and wellbeing, and disclosure appears to have been an important factor in the development of PTG in this study. However, the differences in disclosure may also reflect a lesser need for some mothers to talk about their EM than their SB, in keeping with the differences in CAB and reduced need to reconstruct beliefs following loss in pregnancy.

Individual differences, rather than the nature of the trauma (SB vs EM), were expected to determine rumination. However, significant differences in IR and DR were found between the two groups and may be the result of the differences in the visibility of the pregnancy, time spent pregnant, and expectations women hold of experiencing SB compared to EM. On average, mothers who had experienced an EM revealed higher levels of DR than IR. This is consistent with reports that intrusive thoughts are likely to occur in the immediate aftermath of an event, with deliberate thoughts occurring after the initial shock and distress subside (Black & Wright, 2012). It is therefore interesting that levels of IR and DR were comparable in mothers two to six years after experiencing a SB, which is a novel finding, and may relate to the physical trauma of SB.

### **Strengths and Limitations**

A major strength of this study is the use of a between-groups design to investigate PTG in two groups of mothers following pregnancy loss (SB/EM), overcoming the limitations of previous correlational studies. Another strength is the application of theoretically-derived variables of the Model of Growth in Grief, the first of its kind in

relation to pregnancy loss. The findings provide support for the relevance of the variables in the Model of Growth in Grief (Calhoun et al., 2010) in the development of PTG in mothers following SB and EM. Finally, this study had a well-powered sample and used validated measures to capture key variables identified in the model.

This study has some limitations. In the absence of high-quality validated measures of actual disclosure (a problem encountered in other studies of PTG – see Pietruch and Jobson (2012)), a single-item measure of actual self-disclosure was created for this study. Single-item measures may not be sufficient to accurately measure a construct; however, in the design of this measure, attempts were made to help structure recall of time spent talking about loss. The large variability in responses may demonstrate problems with reliability and validity, but may also be an accurate reflection of individual differences in actual self-disclosure. The DTQ (Müller et al., 2000) was used to aid further understanding of disclosure; however, future research would benefit from developing a validated measure of disclosure to further test the Model of Growth in Grief.

As the majority of participants were White British who were married/civil partnership/cohabiting and educated to A-level or a degree qualification, the transferability of findings to other populations may be limited. Women who do not experience distress following pregnancy loss may not be as motivated to seek opportunities to participate in research. As the sample was self-selecting, it is likely that women who are interested in reflecting on their experience were more likely to participate. Online samples may therefore not be representative and results should be interpreted with caution. However, this is not a great concern here as 98.7% of people aged 16-44 have access to the internet (Office of National Statistics, 2016) and the sample reflected this age group.

The proportion of variance (46.4%) in PTG accounted for in this study indicates there are other factors contributing to its occurrence in mothers two to six years following SB and EM. The model suggests that the development of PTG will take time (involving a process of disclosure and rumination) and although this study replicated the 2-6 year time point used previously by Büchi et al. (2007), there is a lack of understanding in the literature of how time affects PTG and when levels may peak. Having a child since experiencing a pregnancy loss may also impact the development of PTG by potentially affecting the disclosure and rumination aspects of the Model of Growth in Grief. Time since loss and having a child since loss were not included in the regression model as they were not found to be significantly correlated with PTG. However, it is possible that these variables may help to explain a small proportion of the unaccounted variance in PTG. The current study's findings therefore raise further questions about which factors underpin PTG

following SB and EM and future research would benefit from exploring this. Finally, a limited number of confounding variables need to be considered. A greater percentage of mothers following EM had also experienced a LM (14.3%) compared to mothers following SB (1.8%) and the EM group were slightly older than the SB group.

### **Clinical Implications**

Interventions targeting the key variables in the Model of Growth in Grief (CAB, disclosure and rumination) are likely to be clinically useful to promote psychological adjustment in mothers who have experienced SB and EM. The findings are therefore relevant to all practitioners who work with women following loss in pregnancy. In light of the finding that higher PTG and PTSD symptoms were found in mothers following SB than EM, it may be helpful to focus more resources on mothers who experience this type of loss. However, it is important to note that not all loss may result in PTG and some bereaved mothers may find the concept of growth offensive (Tedeschi & Calhoun, 2004a). Whilst PTG is common following trauma, it is not universal, and practitioners should not hold expectations that every mother will experience growth or that it is necessary for psychological adjustment to occur. Where mothers' beliefs have been seriously challenged, clinicians could work to support mothers to talk about their feelings about losing their baby and encourage greater disclosure with friends and family. Ultimately, there remains a need to change attitudes to pregnancy loss (especially EM) and disclosure at a societal level, rather than simply supporting mothers and their families through this type of loss.

### **Conclusions**

Significantly higher levels of PTG, PTSD symptoms and perinatal grief were found in mothers following SB compared to EM. Mothers experienced greater challenges to their assumptive beliefs and revealed higher levels disclosure following their SB. These findings can partially be explained by differences in key variables from the Model of Growth in Grief.

## References

- Adeyemi, A., Mosaku, K., Ajenifuja, O., Fatoye, F., Makinde, N., & Ola, B. (2008). Depressive Symptoms in a Sample of Women Following Perinatal Loss. *Journal of the National Medical Association*, 100(12), 1463-1468. doi:10.1016/S0027-9684(15)31548-0
- Ashbaugh, A. R., Houle-Johnson, S., Herbert, C., El-Hage, W., & Brunet, A. (2016). Psychometric validation of the English and French versions of the Posttraumatic Stress Disorder Checklist for DSM-5 (PCL-5). *PloS one*, 11(10), e0161645.
- Badenhorst, W., & Hughes, P. (2007). Psychological aspects of perinatal loss. *Best Practice & Research Clinical Obstetrics & Gynaecology*, 21(2), 249-259.
- Bellizzi, K. M. (2004). Expressions of generativity and posttraumatic growth in adult cancer survivors. *The International Journal of Aging and Human Development*, 58(4), 267-287.
- Bennett, S. M., Litz, B. T., Lee, B. S., & Maguen, S. (2005). The Scope and Impact of Perinatal Loss: Current Status and Future Directions. *Professional Psychology: Research and Practice*, 36(2), 180.
- Black, B., & Sandelowski, M. (2010). Personal growth after severe fetal diagnosis. *Western Journal of Nursing Research*, 32(8), 1011-1030.
- Black, B. P., & Wright, P. (2012). Posttraumatic growth and transformation as outcomes of perinatal loss. *Illness, Crisis & Loss*, 20(3), 225-237.
- Bonanno, G. A., Wortman, C. B., Lehman, D. R., Tweed, R. G., Haring, M., Sonnega, J., . . . Nesse, R. M. (2002). Resilience to loss and chronic grief: a prospective study from preloss to 18-months postloss. *Journal of personality and social psychology*, 83(5), 1150.
- Brier, N. (2008). Grief following miscarriage: a comprehensive review of the literature. *Journal of Women's Health*, 17(3), 451-464.
- Brownlee, K., & Oikonen, J. (2004). Toward a theoretical framework for perinatal bereavement. *British Journal of Social Work*, 34(4), 517-529.
- Büchi, S., Mörgeli, H., Schnyder, U., Jenewein, J., Glaser, A., Fauchère, J.-C., . . . Sensky, T. (2009). Shared or discordant grief in couples 2–6 years after the death of their



- premature baby: Effects on suffering and posttraumatic growth. *Psychosomatics*, 50(2), 123-130.
- Büchi, S., Mörgeli, H., Schnyder, U., Jenewein, J., Hepp, U., Jina, E., . . . Sensky, T. (2007). Grief and post-traumatic growth in parents 2–6 years after the death of their extremely premature baby. *Psychotherapy and psychosomatics*, 76(2), 106-114.
- Cacciatore, J. (2010). The unique experiences of women and their families after the death of a baby. *Social work in health care*, 49(2), 134-148.
- Cacciatore, J. (2013). *Psychological effects of stillbirth*. Paper presented at the Seminars in Fetal and Neonatal Medicine.
- Cacciatore, J., & Bushfield, S. (2007). Stillbirth: the mother's experience and implications for improving care. *Journal of social work in end-of-life & palliative care*, 3(3), 59-79.
- Cacciatore, J., Rådestad, I., & Frederik Frøen, J. (2008). Effects of contact with stillborn babies on maternal anxiety and depression. *Birth*, 35(4), 313-320.
- Calhoun, L. G., & Tedeschi, R. G. (1990). Positive aspects of critical life problems: Recollections of grief. *Omega-Journal of Death and Dying*, 20(4), 265-272.
- Calhoun, L. G., & Tedeschi, R. G. (2001). Posttraumatic growth: The positive lessons of loss. In R. A. Neimeyer (Ed.), *Meaning reconstruction and the experience of loss* (pp. 157-172). Washington, DC: American Psychological Association.
- Calhoun, L. G., & Tedeschi, R. G. (2006). *The handbook of posttraumatic growth: Research and practice*. Mahwah, NJ: Lawrence Erlbaum Associates Publishers.
- Calhoun, L. G., Tedeschi, R. G., Cann, A., & Hanks, E. A. (2010). Positive outcomes following bereavement: Paths to posttraumatic growth. *Psychologica Belgica*, 50(1-2), 1-2.
- Campbell-Jackson, L., & Horsch, A. (2014). The psychological impact of stillbirth on women: A systematic review. *Illness, Crisis & Loss*, 22(3), 237-256.
- Cann, A., Calhoun, L. G., Tedeschi, R. G., Kilmer, R. P., Gil-Rivas, V., Vishnevsky, T., & Danhauer, S. C. (2010). The Core Beliefs Inventory: A brief measure of disruption in the assumptive world. *Anxiety, Stress & Coping*, 23(1), 19-34.

- Cann, A., Calhoun, L. G., Tedeschi, R. G., Triplett, K. N., Vishnevsky, T., & Lindstrom, C. M. (2011). Assessing posttraumatic cognitive processes: The event related rumination inventory. *Anxiety, Stress, & Coping*, 24(2), 137-156.
- Clauss, D. K. (2009). *Psychological distress following miscarriage and stillbirth: an examination of grief, depression and anxiety in relation to gestational length, women's attributions, perception of care and provision of information*. (pHD thesis), Drexel University.
- Conway, K., & Russell, G. (2000). Couples' grief and experience of support in the aftermath of miscarriage. *Psychology and Psychotherapy: Theory, Research and Practice*, 73(4), 531-545.
- Crawley, R., Lomax, S., & Ayers, S. (2013). Recovering from stillbirth: the effects of making and sharing memories on maternal mental health. *Journal of Reproductive and Infant Psychology*, 31(2), 195-207.
- Cryder, C. H., Kilmer, R. P., Tedeschi, R. G., & Calhoun, L. G. (2006). An exploratory study of posttraumatic growth in children following a natural disaster. *American Journal of Orthopsychiatry*, 76(1), 65-69.
- Cuisinier, M., Kuijpers, J., Hoogduin, C., De Graauw, C., & Janssen, H. (1993). Miscarriage and stillbirth: time since the loss, grief intensity and satisfaction with care. *European Journal of Obstetrics and Gynecology and Reproductive Biology*, 52(3), 163-168.
- Engelhard, I. M., van den Hout, M. A., & Arntz, A. (2001). Posttraumatic stress disorder after pregnancy loss. *General Hospital Psychiatry*, 23(2), 62-66.
- Engelkemeyer, S. M., & Marwit, S. J. (2008). Posttraumatic growth in bereaved parents. *Journal of traumatic stress*, 21(3), 344-346.
- Frøen, J. F., Cacciatore, J., McClure, E. M., Kuti, O., Jokhio, A. H., Islam, M., . . . Committee, L. s. S. S. S. (2011). Stillbirths: why they matter. *The Lancet*, 377(9774), 1353-1366.
- Gold, K. J., Sen, A., & Hayward, R. A. (2010). Marriage and cohabitation outcomes after pregnancy loss. *Pediatrics*, 125, 1202-1207.

- Goldbach, K. R., Dunn, D. S., Toedter, L. J., & Lasker, J. N. (1991). The effects of gestational age and gender on grief after pregnancy loss. *American Journal of Orthopsychiatry*, 61(3), 461.
- Goutaudier, N., Nahi, H., Boudoukha, A., Séjourné, N., & Chabrol, H. (2017). Religious beliefs and post-traumatic growth following stillbirth in a sample Moroccan women. *European Psychiatry*, 41, S724.
- Harms, L., & Talbot, M. (2007). The aftermath of road trauma: Survivors' perceptions of trauma and growth. *Health & Social Work*, 32(2), 129-137.
- Horsch, A., Jacobs, I., & McKenzie-McHarg, K. (2015). Cognitive predictors and risk factors of PTSD following stillbirth: a short-term longitudinal study. *Journal of traumatic stress*, 28(2), 110-117.
- Hughes, P., & Riches, S. (2003). Psychological aspects of perinatal loss. *Current Opinion in Obstetrics and Gynecology*, 15(2), 107-111.
- Isguder, C. K., Batmaz, S., Yildiz, M., Songur, E., Dogru, H. Y., Delibas, I. B., . . . Ozsoy, A. Z. (2017). Relationship Between Miscarriage and Dysfunctional Cognitions About Trauma, Coping Mechanisms, and Posttraumatic Growth. *Journal of Rational-Emotive & Cognitive-Behavior Therapy*, 1-20.
- Janoff-Bulman, R. (2006). Schema-change perspectives on posttraumatic growth. W: LG Calhoun, RG Tedeschi (eds.), *Handbook of posttraumatic growth. Research and practice* (81-99): Mahwah: Lawrence Erlbaum.
- Kersting, A., & Wagner, B. (2012). Complicated grief after perinatal loss. *Dialogues in clinical neuroscience*, 14(2), 187.
- Klier, C. M., Geller, P., & Ritsher, J. (2002). Affective disorders in the aftermath of miscarriage: a comprehensive review. *Archives of Women's Mental Health*, 5(4), 129-149.
- Kluger-Bell, K. (2000). Unspeakable losses: Healing from miscarriage, abortion, and other pregnancy loss.
- Kroenke, K., & Spitzer, R. L. (2002). The PHQ-9: a new depression diagnostic and severity measure. *Psychiatric Annals*, 32(9), 509-515.

- Kroenke, K., Spitzer, R. L., & Williams, J. B. (2001). The phq-9. *Journal of general internal medicine*, 16(9), 606-613.
- Kroenke, K., Strine, T. W., Spitzer, R. L., Williams, J. B., Berry, J. T., & Mokdad, A. H. (2009). The PHQ-8 as a measure of current depression in the general population. *Journal of Affective Disorders*, 114(1), 163-173.
- Krosch, D. J., & Shakespeare-Finch, J. (2017). Grief, traumatic stress, and posttraumatic growth in women who have experienced pregnancy loss. *Psychological Trauma: Theory, Research, Practice, and Policy*, 9(4), 425.
- Laufer, A., & Solomon, Z. (2006). Posttraumatic symptoms and posttraumatic growth among Israeli youth exposed to terror incidents. *Journal of Social and Clinical Psychology*, 25(4), 429-447.
- Lawn, J. E., Blencowe, H., Waiswa, P., Amouzou, A., Mathers, C., Hogan, D., . . . Calderwood, C. (2016). Stillbirths: rates, risk factors, and acceleration towards 2030. *The Lancet*, 387(10018), 587-603.
- Lee, C., & Slade, P. (1996). Miscarriage as a traumatic event: a review of the literature and new implications for intervention. *Journal of psychosomatic research*, 40(3), 235-244.
- Maguen, S., Vogt, D. S., King, L. A., King, D. W., & Litz, B. T. (2006). Posttraumatic growth among Gulf War I veterans: The predictive role of deployment-related experiences and background characteristics. *Journal of Loss and Trauma*, 11(5), 373-388.
- Martin, J. A., Kochanek, K. D., Strobino, D. M., Guyer, B., & MacDorman, M. F. (2005). Annual summary of vital statistics—2003. *Pediatrics*, 115(3), 619-634.
- Mathews, L. L., & Servaty-Seib, H. L. (2007). Hardiness and grief in a sample of bereaved college students. *Death Studies*, 31(3), 183-204.
- Milam, J. (2006). Posttraumatic growth and HIV disease progression. *Journal of consulting and clinical psychology*, 74(5), 817-827.
- Müller, J., Beauducel, A., Raschka, J., & Maercker, A. (2000). Kommunikationsverhalten nach politischer Haft in der DDR—Entwicklung eines Fragebogens zum Offenlegen der Traumaerfahrungen. *Zeitschrift für Politische Psychologie*, 8(4), 413-427.

- Murphy, S., Shevlin, M., & Elklit, A. (2014). Psychological consequences of pregnancy loss and infant death in a sample of bereaved parents. *Journal of Loss and Trauma*, 19(1), 56-69.
- Office of National Statistics. (2016). Statistical bulletin: Internet users: 2016. Retrieved from <https://www.ons.gov.uk/businessindustryandtrade/itandinternetindustry/bulletins/internetusers/2016>
- Pietruch, M., & Jobson, L. (2012). Posttraumatic growth and recovery in people with first episode psychosis: an investigation into the role of self-disclosure. *Psychosis*, 4(3), 213-223.
- Potvin, L., Lasker, J., & Toedter, L. (1989). Measuring grief: A short version of the Perinatal Grief Scale. *Journal of Psychopathology and Behavioral Assessment*, 11(1), 29-45.
- Robinson, G. E. (2011). Dilemmas related to pregnancy loss. *The Journal of nervous and mental disease*, 199(8), 571-574.
- Rosenbach, C., & Renneberg, B. (2008). Positive change after severe burn injuries. *Journal of Burn Care & Research*, 29(4), 638-643.
- Smith, A. C., & Borgers, S. B. (1989). Parental grief response to perinatal death. *Omega-Journal of Death and Dying*, 19(3), 203-214.
- Spitzer, R. L., Kroenke, K., Williams, J. B., & Löwe, B. (2006). A brief measure for assessing generalized anxiety disorder: the GAD-7. *Archives of internal medicine*, 166(10), 1092-1097.
- Tedeschi, R. G., & Calhoun, L. G. (1996). The Posttraumatic Growth Inventory: Measuring the positive legacy of trauma. *Journal of traumatic stress*, 9(3), 455-471.
- Tedeschi, R. G., & Calhoun, L. G. (2004a). A clinical approach to posttraumatic growth. In P. A. Linley & A. Joseph (Eds.), *Positive psychology in practice* (pp. 405-419). Hoboken, NJ: Wiley.
- Tedeschi, R. G., & Calhoun, L. G. (2004b). " Posttraumatic growth: Conceptual foundations and empirical evidence". *Psychological inquiry*, 15(1), 1-18.

- Tedeschi, R. G., & Calhoun, L. G. (2006). Time of change? The spiritual challenges of bereavement and loss. *Omega-Journal of Death and Dying*, 53(1), 105-116.
- Toedter, L. J., Lasker, J. N., & Alhadeff, J. M. (1988). The Perinatal Grief Scale: development and initial validation. *American Journal of Orthopsychiatry*, 58(3), 435.
- Toedter, L. J., Lasker, J. N., & Janssen, H. J. E. M. (2001). International comparison of studies using the perinatal grief scale: a decade of research on pregnancy loss. *Death Studies*, 25(3), 205-228.
- Turton, P., Hughes, P., Evans, C., & Fainman, D. (2001). Incidence, correlates and predictors of post-traumatic stress disorder in the pregnancy after stillbirth. *The British Journal of Psychiatry*, 178(6), 556-560.
- Weathers, F., Litz, B., Keane, T., Palmieri, P., Marx, B., & Schnurr, P. (2013). The PTSD Checklist for DSM-5 (PCL-5). Scale available from the National Center for PTSD. Boston (MA): National Center for PTSD.
- Wright, M. O. D., Crawford, E., & Sebastian, K. (2007). Positive resolution of childhood sexual abuse experiences: The role of coping, benefit-finding and meaning-making. *Journal of Family Violence*, 22(7), 597-608.
- Wright, P. M. (2010). *Pushing on: A grounded theory study of maternal perinatal bereavement*. Loyola University Chicago. Retrieved from [http://ecommons.luc.edu/luc\\_diss/147](http://ecommons.luc.edu/luc_diss/147)



## **Executive Summary**

Losing an unborn child is a very painful experience for a parent and may cause more intense grief than the loss of a partner or parent (Bonanno et al., 2002). Mothers may experience a variety of losses in pregnancy, including early miscarriage (EM; occurring in the first trimester of pregnancy up to 12 weeks gestation), late miscarriage (LM; occurring during the second trimester at 13-23 weeks) and stillbirth (SB; defined as intrauterine death after 24 weeks gestation in the United Kingdom (UK)). Miscarriage affects 200,000 couples every year in the UK and 85% of all miscarriages occur in the first trimester. SB is estimated to occur in nearly 1 in 200 pregnancies (Martin, Kochanek, Strobino, Guyer, & MacDorman, 2005). In comparison to other types of child loss, pregnancy loss is often treated as less significant (Frøen et al., 2011) and parents are left feeling isolated and invalidated in their grief.

Losing a baby through miscarriage (Lee & Slade, 1996) or stillbirth (Horsch, Jacobs, & McKenzie-McHarg, 2015) is recognised as a traumatic event (Kersting & Wagner, 2012). Part of the future is lost when a baby dies and bereavement reactions have been reported to be pervasive, powerful and complex (Cacciatore, 2010; Campbell-Jackson & Horsch, 2014). Mothers are often unprepared for the loss and experience a range of psychological reactions, including denial, guilt, anger, grief and feelings of ‘empty arms’ (Adeyemi et al., 2008; Brownlee & Oikonen, 2004; Cacciatore, 2010; Murphy, Shevlin, & Elklit, 2014). Most studies to date have grouped different perinatal losses in their efforts to understand the psychological impact of loss in pregnancy. However, there are a number of factors that are different between EM and SB and studies have highlighted that the psychological impact of these differing losses requires further clarification (Cacciatore & Bushfield, 2007; Klier, Geller, & Ritscher, 2002).

Individuals who face traumatic experiences often experience a number of distressing emotions, including anxiety, sadness, anger, guilt, depression and desire for a different outcome (Tedeschi & Calhoun, 2004). However, through the journey to make sense of the world post-trauma, evidence suggests that many individuals experience psychological change which they consider to be positive. The phenomenon of PTG has been reported in a wide range of individuals who have faced a variety of traumatic circumstances. While the death of a loved one is a devastating and painful experience, coping with bereavement has also been found to provide a context for significant positive change (Calhoun & Tedeschi, 1990; Engelkemeyer & Marwit, 2008; Mathews & Servaty-Seib, 2007). The possibility of PTG in the aftermath of pregnancy loss has received limited attention to date, with a focus historically on the more traumatic and distressing aspects of



this loss. Black and Wright (2012) highlighted the need for “systematic research on different types of perinatal loss” (e.g. EM and SB) to clarify the posttraumatic “responses that these losses engender” in bereaved parents (p. 233).

Calhoun, Tedeschi, Cann, and Hanks (2010) presented a Model of Growth in Grief to understand how losing “a close other” may result in the acknowledgment of a number of positive changes. Deaths that are unexpected, less ‘natural’ and conflict with or disrupt an individual’s assumptions and beliefs about the way the world should work (e.g. the death of a child) often result in more distress and PTG (Tedeschi & Calhoun, 2006). The Model of Growth in Grief is of particular interest when considering loss in pregnancy, as the majority of published findings of positive changes following loss have focused on the loss of a loved one where there has been a longstanding and ongoing relationship. Although mothers form attachments early on in pregnancy (Gold, Sen, & Hayward, 2010), mothers who experience loss in pregnancy do not have the opportunity to create direct life experiences with their baby, unlike in the death of close family members (Kersting & Wagner, 2012). The pathway to PTG may therefore differ to that found in other types of loss.

This study aimed to investigate PTG in mothers two to six years following SB, compared to in mothers following EM. It also aimed to investigate how theoretically-derived variables of the Model of Growth in Grief (challenge to assumptive beliefs, rumination and disclosure) explain unique variance in PTG when key factors are controlled. The study recruited 120 women who had experienced a SB (57) or EM (63) in the last two to six years. Participants completed questionnaires on emotional distress, disclosure and posttraumatic responses in an online survey. Results indicated that mothers who had experienced a SB experienced significantly greater PTG, posttraumatic stress symptoms and perinatal grief than mothers who had experienced an EM. Greater challenge to assumptive beliefs, intrusive and deliberate rumination, greater urge to talk and consequently greater reported actual self-disclosure were also found in mothers following SB compared to EM.

This is the first study we are aware of that explicitly compared PTG in a SB and EM population. It is also the first study to test the Model of Growth in Grief using a non-correlational design. Interventions that target key variables in the Model of Growth in Grief (challenge to assumptive beliefs, disclosure and rumination) are likely to be clinically useful to promote psychological adjustment in mothers who have experienced SB and EM. The findings are therefore relevant to all clinicians who work with mothers following loss in pregnancy. It must be noted however that whilst PTG is common following trauma, it is not universal, and clinicians should not hold expectations that all mothers will experience PTG or that it is required for psychological adjustment to occur. Ultimately, there remains a need

to change attitudes to pregnancy loss (especially EM) and disclosure at a societal level, rather than simply supporting mothers and their families through this type of loss.

## References

- Adeyemi, A., Mosaku, K., Ajenifuja, O., Fatoye, F., Makinde, N., & Ola, B. (2008). Depressive Symptoms in a Sample of Women Following Perinatal Loss. *Journal of the National Medical Association, 100*(12), 1463-1468. doi:10.1016/S0027-9684(15)31548-0
- Black, B. P., & Wright, P. (2012). Posttraumatic growth and transformation as outcomes of perinatal loss. *Illness, Crisis & Loss, 20*(3), 225-237.
- Bonanno, G. A., Wortman, C. B., Lehman, D. R., Tweed, R. G., Haring, M., Sonnega, J., . . . Nesse, R. M. (2002). Resilience to loss and chronic grief: a prospective study from preloss to 18-months postloss. *Journal of personality and social psychology, 83*(5), 1150.
- Brownlee, K., & Oikonen, J. (2004). Toward a theoretical framework for perinatal bereavement. *British Journal of Social Work, 34*(4), 517-529.
- Cacciatore, J. (2010). The unique experiences of women and their families after the death of a baby. *Social work in health care, 49*(2), 134-148.
- Cacciatore, J., & Bushfield, S. (2007). Stillbirth: the mother's experience and implications for improving care. *Journal of social work in end-of-life & palliative care, 3*(3), 59-79.
- Calhoun, L. G., & Tedeschi, R. G. (1990). Positive aspects of critical life problems: Recollections of grief. *Omega-Journal of Death and Dying, 20*(4), 265-272.
- Calhoun, L. G., Tedeschi, R. G., Cann, A., & Hanks, E. A. (2010). Positive outcomes following bereavement: Paths to posttraumatic growth. *Psychologica Belgica, 50*(1-2), 1-2.
- Campbell-Jackson, L., & Horsch, A. (2014). The psychological impact of stillbirth on women: A systematic review. *Illness, Crisis & Loss, 22*(3), 237-256.
- Engelkemeyer, S. M., & Marwit, S. J. (2008). Posttraumatic growth in bereaved parents. *Journal of traumatic stress, 21*(3), 344-346.

- Frøen, J. F., Cacciatore, J., McClure, E. M., Kuti, O., Jokhio, A. H., Islam, M., . . . Committee, L. s. S. S. S. (2011). Stillbirths: why they matter. *The Lancet*, 377(9774), 1353-1366.
- Gold, K. J., Sen, A., & Hayward, R. A. (2010). Marriage and cohabitation outcomes after pregnancy loss. *Pediatrics*, 125, 1202-1207.
- Horsch, A., Jacobs, I., & McKenzie-McHarg, K. (2015). Cognitive predictors and risk factors of PTSD following stillbirth: a short-term longitudinal study. *Journal of traumatic stress*, 28(2), 110-117.
- Kersting, A., & Wagner, B. (2012). Complicated grief after perinatal loss. *Dialogues in clinical neuroscience*, 14(2), 187.
- Klier, C. M., Geller, P., & Ritsher, J. (2002). Affective disorders in the aftermath of miscarriage: a comprehensive review. *Archives of Women's Mental Health*, 5(4), 129-149.
- Lee, C., & Slade, P. (1996). Miscarriage as a traumatic event: a review of the literature and new implications for intervention. *Journal of psychosomatic research*, 40(3), 235-244.
- Martin, J. A., Kochanek, K. D., Strobino, D. M., Guyer, B., & MacDorman, M. F. (2005). Annual summary of vital statistics—2003. *Pediatrics*, 115(3), 619-634.
- Mathews, L. L., & Servaty-Seib, H. L. (2007). Hardiness and grief in a sample of bereaved college students. *Death Studies*, 31(3), 183-204.
- Murphy, S., Shevlin, M., & Elklit, A. (2014). Psychological consequences of pregnancy loss and infant death in a sample of bereaved parents. *Journal of Loss and Trauma*, 19(1), 56-69.
- Tedeschi, R. G., & Calhoun, L. G. (2004). " Posttraumatic growth: Conceptual foundations and empirical evidence". *Psychological inquiry*, 15(1), 1-18.
- Tedeschi, R. G., & Calhoun, L. G. (2006). Time of change? The spiritual challenges of bereavement and loss. *Omega-Journal of Death and Dying*, 53(1), 105-116.

## **Connecting Narrative**

### **Overview**

This connecting narrative aims to integrate the research projects that have been completed as part of the DClinPsy at the University of Bath. Undertaking these projects has been one of the most enjoyable parts of training. The narrative will first consider the case studies completed on the course before discussing and reflecting on the broader themes characterised by the service improvement project, main research project and critical review of literature.

### **Case studies**

Over the course of clinical training, I completed five case studies across my working age adult, older adult, child, learning disability and adult health placements. The case studies provided valuable research skills and enabled me to draw on theory to make clinically relevant conclusions. Four of the five case studies were cognitive behavioural in their approach, but also drew upon or considered other therapeutic approaches (e.g. third wave CBT/systemic). One case study was written purely from a systemic perspective and provided a welcome lens into alternative ways of formulating with clients and the systems supporting them. Across all case studies, the importance of the therapeutic relationship was highlighted and this theme connects the case studies to my service improvement project.

My first case study highlighted the benefit of including timeline generation in an extended CBT formulation to provide an effective base from which to work on Body Dysmorphic Disorder (BDD) using CBT. Having worked prior to training in an Improving Access to Psychological Therapy (IAPT) service, I was confident in my ability to deliver CBT. However, I was soon struck by the severe and enduring nature of the difficulties I was working with. Working with clients in a secondary care mental health service proved very different and significantly more challenging than anything I had done previously. My first case study therefore taught me the importance of the therapeutic relationship and interventions being client-led. Developing a timeline together and making sense of the past in light of a BDD diagnosis was an important first step and helped to build the therapeutic relationship. Although it would have been helpful to utilise a measure examining the therapeutic relationship directly, the use of idiosyncratic measures throughout the therapy revealed improvements that were not captured by more formal outcome measures. This case study was accepted as a poster at the British Association for Behavioural and Cognitive Psychotherapy 2016 conference in Belfast.

The second case study was completed on my older adult placement in a community-based later life therapies team. I integrated a compassion-focussed approach with CBT to treat low self-esteem and depression experienced by an older woman in the context of physical health difficulties and anxiety about the future. A single case experimental design (SCED) was adopted to evaluate the effectiveness of therapy. This was my first experience of using a SCED and taught me the benefits of using session-by-session measures to monitor progress and outcomes of therapy. Once again, making sense of the past (including consideration of early life experiences and intergenerational beliefs) was an important part of the therapy and lack of social contact/support was identified as an important factor in the maintenance of my client's difficulties. I embraced the slower pace of the work and enjoyed the space to reflect and freedom that this brought in therapy.

My third case study also highlighted the therapeutic relationship as important and focused on supporting an adolescent with chronic headaches to learn to take steps to live alongside her pain. Anxiety-specific CBT was introduced first before using the three pillars of Acceptance and Commitment Therapy (ACT) as a way of engaging in flexible and persistent patterns of values-directed behaviour, whilst in contact with continuing pain. A SCED was used once more to evaluate the intervention and I enjoyed the opportunity to learn more about ACT. Systemic factors were identified as important in this case, which also played a key role on my next placement in a community team for people with learning disabilities.

The fourth case study adopted a systemic approach with a residential staff team to support a woman with Down's Syndrome and moderate intellectual disability experiencing bereavement. This taught me about the importance of working with the systems around people with learning disabilities and raised questions about stigma and discrimination. Developing a trusting and therapeutic relationship with the staff team opened a space for more positive stories about the resident to develop. Staff were left feeling empowered and resourceful and reflecting upon unique outcomes helped to break up the problem-saturated story. I enjoyed the opportunity to work in a different way on this placement and will take this learning forward into my career as a qualified Clinical Psychologist.

The fifth case study explored the utility of transdiagnostic CBT with a patient presenting with generalised anxiety, depression and health anxiety in the context of chronic kidney disease. The case shared similarities with my child and older adult case studies, where psychological distress was also associated with physical health difficulties. A SCED was adopted and completing outcome measures each week enabled progress to be more accurately tracked, including which transdiagnostic interventions resulted in change. Once

again, the therapeutic relationship was an important factor in the patient's willingness to try new ideas.

### **Service Improvement Project**

I first came across my service improvement project (volunteers' experiences of helping hoarders and hoarders' experiences of being helped) at the research fair and it turned out to be the largest project that I completed on clinical training. I was excited about the prospect of working with a charity rather than service in the NHS, as I am aware of the lack of support that charities often have when evaluating and developing their services. I also had an interest in understanding more about the charity's Making Space project and the positive outcomes that had been reported. The service improvement project began with a meeting with the charity's Director of Development (Dan Lyus), Making Space project coordinator (Nia Hall) and my internal supervisor (Dr James Gregory). Together we identified the key aims of the service improvement project and developed a plan to conduct 1:1 interviews with volunteers and clients from the Making Space project.

As the Making Space project was part of a charity, NHS ethical approval was not required and all project documentation was approved by the University of Bath Ethics committee in June 2016. Eleven interviews with clients and volunteers were conducted over the following 18 months. The period of recruitment was significantly longer than anticipated as we faced challenges recruiting clients. Unfortunately, the charity did not have permission to contact a number of previous clients and therefore recruitment had to allow for new clients to be seen by volunteers before they could be interviewed. This experience taught me the importance of patience and flexibility when conducting research! However, in spite of the challenges recruiting clients, I really enjoyed conducting the interviews and meeting the clients and their volunteers. It was a privilege to be invited into clients' homes and hear their stories of how they had developed a problem accumulating and discarding numerous possessions, as well as the difference that the Making Space project had made to their lives. My previous experience of conducting interviews (including qualitative analysis) was invaluable and I enjoyed becoming immersed in the interview data. It was also helpful for a second supervisor, Dr Vuokko Wallace, to join the project prior to the analysis phase and she provided valuable input into the qualitative analysis.

A number of themes were developed and resulted in specific recommendations for the charity, including the need to provide more support for volunteers. The relationship that formed between client and volunteer was crucial in providing a trusting foundation from which clients felt able to move forward. This theme echoed my experiences on placement,

where I have found the therapeutic relationship to be a hugely important part of my work with clients. A peer support group for volunteers has since been developed and it has been really pleasing to see the charity include a number of quotes from clients and volunteers in their project information and applications for additional funding. Being involved in this project really highlighted the difference that third sector organisations can make and it was a privilege to work alongside the charity as they work to grow the Making Space project for the future.

## **Main Research Project and Critical Review of Literature**

As a scientist-practitioner, I believe it is important to actively contribute to the evidence-base and for the evidence-base to inform my clinical practice. I chose to focus most of my research on training in the field of clinical health as this is where the majority of my experience prior to clinical training had been. I was fortunate to work previously on a national project exploring maternal contact with the stillborn infant and developed a number of contacts in perinatal health. I also worked on a project in the Paediatric Intensive Care Unit (PICU) at Bristol Children's Hospital that saw me come into contact with most wards within the hospital. It was from these experiences that the ideas for my main research project and critical review of literature developed. The two projects share the theme of physical health and adjustment.

### **Critical Review of Literature**

In a meeting with my clinical tutor, Dr Cara Davis, I was reminded of some of the difficulties I had come into contact with during my 2.5 years working at Bristol Children's Hospital. Part of my role had involved visiting the cardiac ward to recruit families who had been admitted to PICU following life-saving cardiac treatment. I had been struck by the reactions of patients and their families to their cardiac scar, with a huge variety of responses from disgust to pride. I was keen to find out more about psychological reactions to scarring following cardiac surgery but soon found that there had been very little research to date. Even after broadening the literature search to include other types of planned surgery, it was agreed that there were too few papers to conduct a systematic literature review or meta-analysis.

I began to explore the vast field of visible difference which revealed a wealth of literature in relation to psychological adjustment where disfigurement already existed or where it had occurred through injury. It appeared that less was known about adjustment when treatment had been planned (e.g. surgery). A meeting with my internal supervisors (Professor Paul Salkovskis and Dr Cara Davis) led to the decision to write a conceptual

review that incorporated the existing literature on scarring with research from the visible difference literature to develop a new model to understand psychological adjustment to scarring following planned surgery.

The process of writing the conceptual review was incredibly daunting and challenging as I had not embarked on a project like this before. It was also unclear what the process of writing this type of review would involve and I had to embrace a position of uncertainty, something I have never enjoyed doing! However, reassurance and support from my two internal supervisors and Dr Nell Ellison (Clinical Psychologist for Paediatric Cardiology Services) enabled me to enjoy the process of generating new theoretical knowledge. I found the process of developing a new model exciting and found it particularly rewarding to develop a number of practical recommendations for health professionals working in physical health settings. My previous experience at Bristol Children's Hospital highlighted the importance of preparing and supporting patients through their surgery and resultant scarring and I was pleased that this review offered some suggestions for how multidisciplinary teams might do this. Writing the conceptual review was inspired by my experience in health prior to training, similar to my main research project.

### **Main Research Project**

I was keen for my main research project to focus on the area of perinatal health as this is an area I have always been interested in. I was already in contact with a past research supervisor, Professor Antje Horsch, who is very experienced in conducting research and working clinically with women who have experienced perinatal loss. We had worked together prior to clinical training on a large scale national project interviewing women who had experienced a stillbirth. This experience proved invaluable as I sought to immerse myself in the area of perinatal loss once again.

Through a number of skype meetings with my internal supervisor (Dr Megan Wilkinson-Tough) and two external supervisors (Professor Antje Horsch and Dr Sarah Stacey), we agreed to investigate emotional reactions to loss in pregnancy. I was particularly interested in understanding more about the positive changes that might occur as a result of loss in pregnancy, as this was a theme that had emerged from the interviews I had conducted with mothers previously. However, settling on the two comparison groups proved very challenging, as it did not feel appropriate to use a control group of women who had not experienced pregnancy loss, nor to compare pregnancy loss to another type of bereavement. In light of my interest in how disclosure may be involved in the development



of posttraumatic growth, I chose to investigate posttraumatic growth in mothers who had experienced an early miscarriage and mothers who had experienced a stillbirth.

It was important to include people with personal experience in the design of the project, especially as the project was going to be hosted online and therefore involve no direct contact with the research team. The project replicated the time point used in a previous study on perinatal loss and therefore meant that recruiting through the NHS would not be possible as local services did not have permission to retrospectively contact patients. I knew that the process of obtaining IRAS approval was often confusing, slow and frustrating, so I was relieved that I did not have to go through IRAS to recruit through the NHS. However, in hindsight, I wonder if this was a missed opportunity for valuable learning and practice for the future. As I intend to continue conducting research in my qualified practice, I look forward to learning this skill.

I met early on with a couple who had experienced a stillbirth and later a woman who had experienced several early miscarriages. Both meetings highlighted how vital service user consultation is and provided valuable feedback on the project and associated documentation. The Stillbirth and Neonatal Death Charity (Sands) were consulted regarding study documentation and agreed to support and advertise the project. Although I had some apprehension about how mothers might interpret questions relating to positive changes that they may have experienced following their loss, this did not arise as an issue at any stage of the project. The greatest surprise and challenge was instead the backlash that came from bereaved parents who did not meet the eligibility criteria for the study. I was aware of the sensitivity surrounding loss in pregnancy but was not prepared for the intensity nor negativity of the comments that were posted on the Sands Facebook page in the hours after they first posted information about the project. This served as a valuable learning opportunity for me and for Sands, and highlighted the lack of awareness in the general population about the need for inclusion and exclusion criteria in research studies. It also brought home the reality of using social media to advertise research and the negative consequences of instant messaging. Sands have since developed a webpage explaining why studies need inclusion and exclusion criteria in research and they are in the process of creating a separate page away from their Facebook group to advertise research studies in the future. Subsequent posts by Sands in support of the project did not generate further negative comments and to my surprise, our recruitment target was exceeded as we recruited a total of 120 mothers.

## Summary

I believe my training on the Bath course has put me in an excellent position to continue undertaking research in my future career as a qualified Clinical Psychologist. Undertaking the research components of the course has been one of the most enjoyable aspects of clinical training. I have found conducting research incredibly rewarding and valued the opportunity to explore a number of areas in detail using multiple research methodologies, rather than researching one topic. However, conducting multiple research projects alongside working full-time was stressful and frustrating at times as I felt I could not give my full attention to each project. I hope that my dedicated half day per week for continuous professional development and research in my first qualified post will help to ease this pressure in the future.

Across all the case studies, the importance of the therapeutic relationship was highlighted and this theme connected my case studies to my service improvement project. My main research project and critical review of literature were both conducted in the field of clinical health and shared the theme of physical health and adjustment. Conducting three research projects and five case studies on the course has taught me a great deal about research design and conduct and I believe this variety of research designs and methodologies will stand in me in good stead as I start my next chapter in qualified practice.

Finally, the course has taught me the equal value and importance of undertaking service improvement, large scale research and single case experimental designs in order to grow and develop the evidence base in Clinical Psychology. I am committed to being an evidence-based practitioner who contributes to the evidence base. I am grateful for the variety of experiences I have had on placement to practice new skills, apply theory to practice and write up examples of my clinical work. I have valued the involvement of people with personal experience in my research and the opportunity to write up the findings for publication. I hope that my research will make a valuable contribution to the evidence base and be useful to clinicians in the future.



## Acknowledgments

First and foremost, I would like to express how indebted I am to my husband Tim for his unwavering support through the ups and downs of clinical training. His love, patience and constant belief in me has given me the motivation to keep going. I hope I can one day repay all the support he has given me over the last three years. I would also like to thank my family, friends and fellow trainees for all their support through clinical training.

I wish to thank Dr Cara Davis, my Clinical Tutor, for overseeing my professional and personal development and generally being a huge support in time of need throughout the course. I would also like to thank all of my placement supervisors (Dr Kerry Ozcelik, Dr Hen Joannidi, Dr Rebecca Guhan, Dr Chloe Constable, Dr Samantha Green, Dr Suzanne Whitehead, Dr Kate Druett, Dr Leon Dysch and Marion Dixon) for their support in completing my case studies and developing my clinical skills. Working alongside these inspiring supervisors has motivated me to be the best Clinical Psychologist I can be.

I have been fortunate to have four incredibly supportive and passionate academic supervisors for my three research projects on clinical training. I am grateful to Professor Paul Salkovskis, who supported me through the daunting and challenging journey of writing a conceptual literature review. I would also like to thank Dr Cara Davis and Dr Nell Ellison for their encouragement and involvement in the review. I am grateful to Dr James Gregory for the opportunity to conduct a service improvement project in a third sector organisation and to further refine my skills in qualitative methodology. I would also like to thank Dr Megan Wilkinson-Tough, Professor Antje Horsch and Dr Sarah Stacey for their support in continuing my interest in perinatal health. In spite of the challenges that were faced along the way, this was my favourite project on training. I hope that my findings will make a real difference to the lives of mothers and their families who have experienced loss in pregnancy in the future.

Finally, I would like to thank all the clients that I had the privilege to work alongside and participants who agreed to take part in my research. Without their involvement, I wouldn't be where I am today.

Kirsty Ryninks

University of Bath

May 2018



## Appendices

### Appendix I: Instructions for Authors – Body Image



## BODY IMAGE

An International Journal of Research

### AUTHOR INFORMATION PACK

#### TABLE OF CONTENTS

• Description	p.1
• Impact Factor	p.2
• Abstracting and Indexing	p.2
• Editorial Board	p.2
• Guide for Authors	p.4



ISSN: 1740-1445

#### DESCRIPTION

*Body Image* is an international, peer-reviewed journal that publishes high-quality, scientific articles on **body image** and human **physical appearance**. Body image is a multi-faceted concept that refers to persons' perceptions and attitudes about their own body, particularly but not exclusively its appearance. The journal invites contributions from a broad range of disciplines - psychological science, other social and behavioral sciences, and medical and health sciences. The journal publishes original research articles, brief research reports, theoretical and review papers, and science-based practitioner reports of interest. The journal gives an annual award for the best doctoral dissertation in this field.

Suitable topics for [submission](#) of manuscripts include:

- The effects of body image and physical characteristics (e.g., body size, attractiveness, physical disfigurements or disorders) on **psychological functioning**, interpersonal processes, and quality of life;
- Body image and physical appearance in the full range of medical and allied health contexts;
- Body image and physical appearance in diverse cultural contexts;
- Validation of assessments of the multidimensional body image construct;
- Factors that influence positive and negative body image development;
- Adaptive and maladaptive body image processes and their clinically relevant consequences on **psychosocial** functioning and quality of life;
- Relationship of body image to **behavioral** variables (e.g., exercise and other physical activity, eating and weight-control behaviors, grooming and appearance-modifying behaviors, and social behaviors);
- Scientific evaluation of interventions to promote positive body image or to prevent or treat body image difficulties and disorders.

#### Benefits to authors

We also provide many author benefits, such as free PDFs, a liberal copyright policy, special discounts on Elsevier publications and much more. Please click [here](#) for more information on our [author services](#).

Please see our [Guide for Authors](#) for information on article submission. If you require any further information or help, please visit our [Support Center](#)

## IMPACT FACTOR

---

2016: 2.926 © Clarivate Analytics Journal Citation Reports 2017

## ABSTRACTING AND INDEXING

---

Scopus  
CINAHL  
PsycINFO  
Current Contents/Social & Behavioral Sciences  
Google Scholar  
MEDLINE®

## EDITORIAL BOARD

---

### *Editor-in-Chief*

**Tracy L. Tylka**, Dept. of Psychology, The Ohio State University, 1465 Mt. Vernon Avenue, Marion, Ohio, 43302, USA, Fax: +1 614 292 5817

### *Founding Editor*

**Thomas F. Cash**

### *Associate Editors*

**Jessica M. Alleva**, Maastricht University, Maastricht, Netherlands  
**Rachel M. Calogero**, Western University, London, Ontario, Canada  
**Matthew Fuller-Tyszkiewicz**, Deakin University, Burwood, Victoria, Australia  
**Sarah K. Murnen**, Kenyon College, Gambier, Ohio, USA  
**Rachel F. Rodgers**, Northeastern University, Boston, Massachusetts, USA  
**Viren Swami**, Anglia Ruskin University, Cambridge, UK

### *Editorial Board Members*

**Carolyn B. Becker**, Trinity University, San Antonio, Texas, USA  
**Kristin Billaud Feragen**, Oslo University Hospital, Oslo, Norway  
**Aaron J. Blashill**, Massachusetts General Hospital, Charlestown, Massachusetts, USA  
**Catherine P. Cook-Cottone**, University at Buffalo, Buffalo, New York, USA  
**Alexandra F. Corning**, University of Notre Dame, Notre Dame, Indiana, USA  
**Canice E. Crerand**, Ohio State University Medical Center, Columbus, Ohio, USA  
**Phillippa C. Diedrichs**, University of the West of England, Bristol, UK  
**Jasmine Fardouly**, Macquarie University, Sydney, New South Wales, Australia  
**Helen J. Fawcner**, Leeds Beckett University, Leeds, UK  
**Ellen E. Fitzsimmons-Craft**, Washington University in St. Louis School of Medicine, St. Louis, Missouri, USA  
**Debra L. Franko**, Northeastern University, Boston, Massachusetts, USA  
**David Frederick**, Chapman University, Orange, California, USA  
**Rick M. Gardner**, University of Colorado Denver, Denver, Colorado, USA  
**Meghan M. Gillen**, Penn State University, University Park, Pennsylvania, USA  
**David H. Gleaves**, University of South Australia, Adelaide, South Australia, Australia  
**Scott Griffiths**, The University of Melbourne, Victoria, Australia  
**Sarah Grogan**, Manchester Metropolitan University, Manchester, UK  
**Emma Halliwell**, University of the West of England, Bristol, UK  
**Diana Harcourt**, University of the West of England, Bristol, UK  
**Thomas B. Hildebrandt**, Mount Sinai School of Medicine, New York, New York, USA  
**Andrew J. Hill**, University of Leeds, Leeds, UK  
**Kristina Holmqvist Gattario**, Göteborgs Universitet, Göteborg, Sweden  
**Kristin J. Homan**, Grove City College, Grove City, Pennsylvania, USA  
**Todd Jackson**, University of Macau, Macau, China  
**Josee L. Jarry**, University of Windsor, Windsor, Ontario, Canada  
**Bryan T. Karazsia**, The College of Wooster, Wooster, Ohio, USA  
**Allison C. Kelly**, University of Waterloo, Waterloo, Ontario, Canada  
**Ashley Kroon Van Diest**, Cincinnati Children's Hospital, Cincinnati, Ohio, USA  
**Ingela L. Kvale**, University of Oslo, Oslo, Norway  
**James E. Leone**, Bridgewater State University, Bridgewater, Massachusetts, USA  
**Michael P. Levine**, Kenyon College, Gambier, Ohio, USA  
**Janet M. Liechty**, University of Illinois at Urbana-Champaign, Urbana, Illinois, USA  
**Miriam Liss**, University of Mary Washington, Fredericksburg, Virginia, USA

**Leanne Magee**, The Children's Hospital of Philadelphia, Philadelphia, Pennsylvania, USA  
**Charlotte N. Markey**, Rutgers University, Camden, New Jersey, USA  
**Kathleen A. Martin Ginis**, University of British Columbia, Kelowna, British Columbia, Canada  
**Marita McCabe**, Australian Catholic University, Melbourne, Victoria, Australia  
**Don R. McCreary**, Brock University, St. Catharines, Ontario, Canada  
**Todd G. Morrison**, University of Saskatchewan, Saskatoon, Saskatchewan, Canada  
**Kate E. Mulgrew**, University of the Sunshine Coast, Sippy Downs, Queensland, Australia  
**Stuart B. Murray**, University of California at San Francisco (UCSF), San Francisco, California, USA  
**Taryn A. Myers**, Virginia Wesleyan College, Norfolk, Virginia, USA  
**Michael C. Parent**, University of Texas at Austin, Austin, Texas, USA  
**Susan J. Paxton**, La Trobe University, Melbourne (Bundoora), Victoria, Australia  
**Trent A. Petrie**, University of North Texas, Denton, Texas, USA  
**Katharine A. Phillips**, Brown University, Providence, Rhode Island, USA  
**Niva Piran**, University of Toronto, Toronto, Ontario, Canada  
**Lina A. Ricciardelli**, Deakin University, Melbourne, Victoria, Australia  
**David B. Sarwer**, Temple University, Philadelphia, Pennsylvania, USA  
**Lauren M. Schaefer**, Neuropsychiatric Research Institute, Fargo, North Dakota, USA  
**Gemma Sharp**, Monash University, Melbourne, Victoria, Australia  
**Linda Smolak**, Kenyon College, Gambier, Ohio, USA  
**J. Kevin Thompson**, University of South Florida, Tampa, Florida, USA  
**Idia B. Thurston**, University of Memphis, Memphis, Tennessee, USA  
**Marika Tiggemann**, Flinders University, Adelaide, South Australia, Australia  
**Lenny R. Vartanian**, UNSW Australia, Sydney, New South Wales, Australia  
**David Veale**, Maudsley Hospital, London, UK  
**Cortney S. Warren**, University of Nevada, Las Vegas, Nevada, USA  
**Jennifer B. Webb**, University of North Carolina at Charlotte, Charlotte, North Carolina, USA  
**Eleanor H. Wertheim**, La Trobe University, Melbourne (Bundoora), Victoria, Australia  
**Nichole L. Wood-Barcalow**, Chalmers P. Wylie VA Ambulatory Care Center, Columbus, Ohio, USA  
**Zali Yager**, Victoria University, Footscray, Victoria, Australia  
**Yuko Yamamiya**, Temple University, Tokyo, Japan



## GUIDE FOR AUTHORS

---

### **Types of Papers**

The journal publishes original research articles, brief research reports, theoretical and review papers, and science-based practitioner reports of interest. The journal also gives an annual award for the best doctoral dissertation in this field.

**Brief Research Reports.** These should not exceed 2,500 words (excluding abstract, references, tables, figures and appendices). Up to a total of two one-page tables, figures, and/or appendices are permitted. The number of references cannot exceed 25.

While regular-length papers have no explicit limits in terms of numbers of words, tables/figures, and references, authors are encouraged to keep their length below 35 total pages. A paper's length must be justified by its empirical strength and the significance of its contribution to the literature.

### **The Seymour Fisher Outstanding Body Image Dissertation Annual Award**

The journal gives an annual award for the best doctoral dissertation in this field. [Click here](#) for more information.

### **Submission checklist**

You can use this list to carry out a final check of your submission before you send it to the journal for review. Please check the relevant section in this Guide for Authors for more details.

#### **Ensure that the following items are present:**

One author has been designated as the corresponding author with contact details:

- E-mail address
- Full postal address

All necessary files have been uploaded:

*Manuscript:*

- Include keywords
- All figures (include relevant captions)
- All tables (including titles, description, footnotes)
- Ensure all figure and table citations in the text match the files provided
- Indicate clearly if color should be used for any figures in print

*Graphical Abstracts / Highlights files* (where applicable)

*Supplemental files* (where applicable)

Further considerations

- Manuscript has been 'spell checked' and 'grammar checked'
- All references mentioned in the Reference List are cited in the text, and vice versa
- Permission has been obtained for use of copyrighted material from other sources (including the Internet)
- A competing interests statement is provided, even if the authors have no competing interests to declare
- Journal policies detailed in this guide have been reviewed
- Referee suggestions and contact details provided, based on journal requirements

For further information, visit our [Support Center](#).

## **BEFORE YOU BEGIN**

### **Ethics in publishing**

Please see our information pages on [Ethics in publishing](#) and [Ethical guidelines for journal publication](#).

### **Human and animal rights**

If the work involves the use of human subjects, the author should ensure that the work described has been carried out in accordance with [The Code of Ethics of the World Medical Association](#) (Declaration of Helsinki) for experiments involving humans; [Uniform Requirements for manuscripts submitted to](#)

**Biomedical journals.** Authors should include a statement in the manuscript that informed consent was obtained for experimentation with human subjects. The privacy rights of human subjects must always be observed.

All animal experiments should comply with the [ARRIVE guidelines](#) and should be carried out in accordance with the U.K. Animals (Scientific Procedures) Act, 1986 and associated guidelines, [EU Directive 2010/63/EU for animal experiments](#), or the National Institutes of Health guide for the care and use of Laboratory animals (NIH Publications No. 8023, revised 1978) and the authors should clearly indicate in the manuscript that such guidelines have been followed.

#### **Declaration of interest**

All authors must disclose any financial and personal relationships with other people or organizations that could inappropriately influence (bias) their work. Examples of potential conflicts of interest include employment, consultancies, stock ownership, honoraria, paid expert testimony, patent applications/registrations, and grants or other funding. Authors must disclose any interests in two places: 1. A summary declaration of interest statement in the title page file (if double-blind) or the manuscript file (if single-blind). If there are no interests to declare then please state this: 'Declarations of interest: none'. This summary statement will be ultimately published if the article is accepted. 2. Detailed disclosures as part of a separate Declaration of Interest form, which forms part of the journal's official records. It is important for potential interests to be declared in both places and that the information matches. [More information.](#)

#### **Submission declaration and verification**

Submission of an article implies that the work described has not been published previously (except in the form of an abstract, a published lecture or academic thesis, see '[Multiple, redundant or concurrent publication](#)' for more information), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. To verify originality, your article may be checked by the originality detection service [Crossref Similarity Check](#).

#### **Preprints**

Please note that [preprints](#) can be shared anywhere at any time, in line with Elsevier's [sharing policy](#). Sharing your preprints e.g. on a preprint server will not count as prior publication (see '[Multiple, redundant or concurrent publication](#)' for more information).

#### **Changes to authorship**

Authors are expected to consider carefully the list and order of authors **before** submitting their manuscript and provide the definitive list of authors at the time of the original submission. Any addition, deletion or rearrangement of author names in the authorship list should be made only **before** the manuscript has been accepted and only if approved by the journal Editor. To request such a change, the Editor must receive the following from the **corresponding author**: (a) the reason for the change in author list and (b) written confirmation (e-mail, letter) from all authors that they agree with the addition, removal or rearrangement. In the case of addition or removal of authors, this includes confirmation from the author being added or removed.

Only in exceptional circumstances will the Editor consider the addition, deletion or rearrangement of authors **after** the manuscript has been accepted. While the Editor considers the request, publication of the manuscript will be suspended. If the manuscript has already been published in an online issue, any requests approved by the Editor will result in a corrigendum.

#### **Article transfer service**

This journal is part of our Article Transfer Service. This means that if the Editor feels your article is more suitable in one of our other participating journals, then you may be asked to consider transferring the article to one of those. If you agree, your article will be transferred automatically on your behalf with no need to reformat. Please note that your article will be reviewed again by the new journal. [More information.](#)

#### **Copyright**

Upon acceptance of an article, authors will be asked to complete a 'Journal Publishing Agreement' (see [more information](#) on this). An e-mail will be sent to the corresponding author confirming receipt of the manuscript together with a 'Journal Publishing Agreement' form or a link to the online version of this agreement.

Subscribers may reproduce tables of contents or prepare lists of articles including abstracts for internal circulation within their institutions. [Permission](#) of the Publisher is required for resale or distribution outside the institution and for all other derivative works, including compilations and translations. If excerpts from other copyrighted works are included, the author(s) must obtain written permission from the copyright owners and credit the source(s) in the article. Elsevier has [preprinted forms](#) for use by authors in these cases.

For gold open access articles: Upon acceptance of an article, authors will be asked to complete an 'Exclusive License Agreement' ([more information](#)). Permitted third party reuse of gold open access articles is determined by the author's choice of [user license](#).

#### **Author rights**

As an author you (or your employer or institution) have certain rights to reuse your work. [More information](#).

#### *Elsevier supports responsible sharing*

Find out how you can [share your research](#) published in Elsevier journals.

#### **Role of the funding source**

You are requested to identify who provided financial support for the conduct of the research and/or preparation of the article and to briefly describe the role of the sponsor(s), if any, in study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the article for publication. If the funding source(s) had no such involvement then this should be stated.

#### *Funding body agreements and policies*

Elsevier has established a number of agreements with funding bodies which allow authors to comply with their funder's open access policies. Some funding bodies will reimburse the author for the gold open access publication fee. Details of [existing agreements](#) are available online.

#### **Open access**

This journal offers authors a choice in publishing their research:

#### **Subscription**

- Articles are made available to subscribers as well as developing countries and patient groups through our [universal access programs](#).
- No open access publication fee payable by authors.
- The Author is entitled to post the [accepted manuscript](#) in their institution's repository and make this public after an embargo period (known as green Open Access). The [published journal article](#) cannot be shared publicly, for example on ResearchGate or Academia.edu, to ensure the sustainability of peer-reviewed research in journal publications. The embargo period for this journal can be found below.

#### **Gold open access**

- Articles are freely available to both subscribers and the wider public with permitted reuse.
- A gold open access publication fee is payable by authors or on their behalf, e.g. by their research funder or institution.

Regardless of how you choose to publish your article, the journal will apply the same peer review criteria and acceptance standards.

For gold open access articles, permitted third party (re)use is defined by the following [Creative Commons user licenses](#):

#### *Creative Commons Attribution (CC BY)*

Lets others distribute and copy the article, create extracts, abstracts, and other revised versions, adaptations or derivative works of or from an article (such as a translation), include in a collective work (such as an anthology), text or data mine the article, even for commercial purposes, as long as they credit the author(s), do not represent the author as endorsing their adaptation of the article, and do not modify the article in such a way as to damage the author's honor or reputation.



*Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)*

For non-commercial purposes, lets others distribute and copy the article, and to include in a collective work (such as an anthology), as long as they credit the author(s) and provided they do not alter or modify the article.

The gold open access publication fee for this journal is **USD 1800**, excluding taxes. Learn more about Elsevier's pricing policy: <https://www.elsevier.com/openaccesspricing>.

*Green open access*

Authors can share their research in a variety of different ways and Elsevier has a number of green open access options available. We recommend authors see our [green open access page](#) for further information. Authors can also self-archive their manuscripts immediately and enable public access from their institution's repository after an embargo period. This is the version that has been accepted for publication and which typically includes author-incorporated changes suggested during submission, peer review and in editor-author communications. Embargo period: For subscription articles, an appropriate amount of time is needed for journals to deliver value to subscribing customers before an article becomes freely available to the public. This is the embargo period and it begins from the date the article is formally published online in its final and fully citable form. [Find out more](#).

This journal has an embargo period of 24 months.

*Elsevier Researcher Academy*

[Researcher Academy](#) is a free e-learning platform designed to support early and mid-career researchers throughout their research journey. The "Learn" environment at Researcher Academy offers several interactive modules, webinars, downloadable guides and resources to guide you through the process of writing for research and going through peer review. Feel free to use these free resources to improve your submission and navigate the publication process with ease.

*Language (usage and editing services)*

Please write your text in good English (American or British usage is accepted, but not a mixture of these). Authors who feel their English language manuscript may require editing to eliminate possible grammatical or spelling errors and to conform to correct scientific English may wish to use the [English Language Editing service](#) available from Elsevier's WebShop.

**Submission**

Our online submission system guides you stepwise through the process of entering your article details and uploading your files. The system converts your article files to a single PDF file used in the peer-review process. Editable files (e.g., Word, LaTeX) are required to typeset your article for final publication. All correspondence, including notification of the Editor's decision and requests for revision, is sent by e-mail.

**Peer review**

This journal operates a double blind review process. All contributions will be initially assessed by the editor for suitability for the journal. Papers deemed suitable are then typically sent to a minimum of two independent expert reviewers to assess the scientific quality of the paper. The Editor is responsible for the final decision regarding acceptance or rejection of articles. The Editor's decision is final. [More information on types of peer review](#).

**Double-blind review**

This journal uses double-blind review, which means the identities of the authors are concealed from the reviewers, and vice versa. [More information](#) is available on our website. To facilitate this, please include the following separately:

*Title page (with author details):* This should include the title, authors' names, affiliations, acknowledgements and any Declaration of Interest statement, and a complete address for the corresponding author including an e-mail address.

*Blinded manuscript (no author details):* The main body of the paper (including the references, figures, tables and any acknowledgements) should not include any identifying information, such as the authors' names or affiliations.

*Use of word processing software*

It is important that the file be saved in the native format of the word processor used. The text should be in single-column format. Keep the layout of the text as simple as possible. Most formatting codes will be removed and replaced on processing the article. In particular, do not use the word processor's options to justify text or to hyphenate words. However, do use bold face, italics, subscripts, superscripts etc. When preparing tables, if you are using a table grid, use only one grid for each

individual table and not a grid for each row. If no grid is used, use tabs, not spaces, to align columns. The electronic text should be prepared in a way very similar to that of conventional manuscripts (see also the [Guide to Publishing with Elsevier](#)). Note that source files of figures, tables and text graphics will be required whether or not you embed your figures in the text. See also the section on Electronic artwork.

To avoid unnecessary errors you are strongly advised to use the 'spell-check' and 'grammar-check' functions of your word processor.

### Article structure

#### Introduction

State the objectives of the work and provide an adequate background, avoiding a detailed literature survey or a summary of the results.

#### Material and methods

Provide sufficient details to allow the work to be reproduced by an independent researcher. Methods that are already published should be summarized, and indicated by a reference. If quoting directly from a previously published method, use quotation marks and also cite the source. Any modifications to existing methods should also be described.

#### Results

Results should be clear and concise, describing the findings and their associated statistical basis. Consider the use of tables and figures for statistical details.

#### Discussion

This section should present the theoretical, empirical, and applied implications of the results, not simply repeat the findings. The study's limitations should be explicitly recognized. A combined Results and Discussion section may be appropriate.

#### Conclusions

The main conclusions of the study may be presented in a short Conclusions section, which may stand alone or form a subsection of a Discussion or Results and Discussion section.

#### Appendices

If there is more than one appendix, they should be identified as A, B, etc. Formulae and equations in appendices should be given separate numbering: Eq. (A.1), Eq. (A.2), etc.; in a subsequent appendix, Eq. (B.1) and so on. Similarly for tables and figures: Table A.1; Fig. A.1, etc.

### Essential title page information

- **Title.** Concise and informative. Titles are often used in information-retrieval systems. Avoid abbreviations and formulae where possible.
- **Author names and affiliations.** Please clearly indicate the given name(s) and family name(s) of each author and check that all names are accurately spelled. You can add your name between parentheses in your own script behind the English transliteration. Present the authors' affiliation addresses (where the actual work was done) below the names. Indicate all affiliations with a lower-case superscript letter immediately after the author's name and in front of the appropriate address. Provide the full postal address of each affiliation, including the country name and, if available, the e-mail address of each author.
- **Corresponding author.** Clearly indicate who will handle correspondence at all stages of refereeing and publication, also post-publication. This responsibility includes answering any future queries about Methodology and Materials. **Ensure that the e-mail address is given and that contact details are kept up to date by the corresponding author.**
- **Present/permanent address.** If an author has moved since the work described in the article was done, or was visiting at the time, a 'Present address' (or 'Permanent address') may be indicated as a footnote to that author's name. The address at which the author actually did the work must be retained as the main, affiliation address. Superscript Arabic numerals are used for such footnotes.

### Abstract

A concise and factual abstract is required. The abstract should state briefly the purpose of the research, the principal results and major conclusions. An abstract is often presented separately from the article, so it must be able to stand alone. For this reason, References should be avoided, but if essential, then cite the author(s) and year(s). Also, non-standard or uncommon abbreviations should be avoided, but if essential they must be defined at their first mention in the abstract itself.

The abstract should be a maximum of 150 words.

#### *Graphical abstract*

Although a graphical abstract is optional, its use is encouraged as it draws more attention to the online article. The graphical abstract should summarize the contents of the article in a concise, pictorial form designed to capture the attention of a wide readership. Graphical abstracts should be submitted as a separate file in the online submission system. Image size: Please provide an image with a minimum of 531 × 1328 pixels (h × w) or proportionally more. The image should be readable at a size of 5 × 13 cm using a regular screen resolution of 96 dpi. Preferred file types: TIFF, EPS, PDF or MS Office files. You can view [Example Graphical Abstracts](#) on our information site.

Authors can make use of Elsevier's [Illustration Services](#) to ensure the best presentation of their images and in accordance with all technical requirements.

#### *Highlights*

Highlights are mandatory for this journal. They consist of a short collection of bullet points that convey the core findings of the article and should be submitted in a separate editable file in the online submission system. Please use 'Highlights' in the file name and include 3 to 5 bullet points (maximum 85 characters, including spaces, per bullet point). You can view [example Highlights](#) on our information site.

#### **Keywords**

Immediately after the abstract, provide a maximum of 6 keywords, using American spelling and avoiding general and plural terms and multiple concepts (avoid, for example, 'and', 'of'). Be sparing with abbreviations: only abbreviations firmly established in the field may be eligible. These keywords will be used for indexing purposes.

#### **Abbreviations**

For economy, consider using abbreviations or acronyms for key terms that appear often in the paper. Introduce the abbreviation parenthetically after the term's first mention in the paper. Ensure consistency of abbreviations throughout the paper. Such abbreviations that are unavoidable in the abstract must be defined at their first mention there, as well as in the footnote. Ensure consistency of abbreviations throughout the article.

#### *Acknowledgements*

Collate acknowledgements in a separate section at the end of the article before the references and do not, therefore, include them on the title page, as a footnote to the title or otherwise. List here those individuals who provided help during the research (e.g., providing language help, writing assistance or proof reading the article, etc.).

#### *Formatting of funding sources*

List funding sources in this standard way to facilitate compliance to funder's requirements:

Funding: This work was supported by the National Institutes of Health [grant numbers xxxx, yyyy]; the Bill & Melinda Gates Foundation, Seattle, WA [grant number zzzz]; and the United States Institutes of Peace [grant number aaaa].

It is not necessary to include detailed descriptions on the program or type of grants and awards. When funding is from a block grant or other resources available to a university, college, or other research institution, submit the name of the institute or organization that provided the funding.

If no funding has been provided for the research, please include the following sentence:

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

#### *Math formulae*

Please submit math equations as editable text and not as images. Present simple formulae in line with normal text where possible and use the solidus (/) instead of a horizontal line for small fractional terms, e.g., X/Y. In principle, variables are to be presented in italics. Powers of e are often more conveniently denoted by exp. Number consecutively any equations that have to be displayed separately from the text (if referred to explicitly in the text).



#### *Footnotes*

Footnotes should be used sparingly. Number them consecutively throughout the article. Many word processors can build footnotes into the text, and this feature may be used. Otherwise, please indicate the position of footnotes in the text and list the footnotes themselves separately at the end of the article. Do not include footnotes in the Reference list.

#### **Artwork**

##### *Electronic artwork*

##### *General points*

- Make sure you use uniform lettering and sizing of your original artwork.
- Embed the used fonts if the application provides that option.
- Aim to use the following fonts in your illustrations: Arial, Courier, Times New Roman, Symbol, or use fonts that look similar.
- Number the illustrations according to their sequence in the text.
- Use a logical naming convention for your artwork files.
- Provide captions to illustrations separately.
- Size the illustrations close to the desired dimensions of the published version.
- Submit each illustration as a separate file.

A detailed [guide on electronic artwork](#) is available.

**You are urged to visit this site; some excerpts from the detailed information are given here.**

##### *Formats*

If your electronic artwork is created in a Microsoft Office application (Word, PowerPoint, Excel) then please supply 'as is' in the native document format.

Regardless of the application used other than Microsoft Office, when your electronic artwork is finalized, please 'Save as' or convert the images to one of the following formats (note the resolution requirements for line drawings, halftones, and line/halftone combinations given below):

EPS (or PDF): Vector drawings, embed all used fonts.

TIFF (or JPEG): Color or grayscale photographs (halftones), keep to a minimum of 300 dpi.

TIFF (or JPEG): Bitmapped (pure black & white pixels) line drawings, keep to a minimum of 1000 dpi.

TIFF (or JPEG): Combinations bitmapped line/half-tone (color or grayscale), keep to a minimum of 500 dpi.

##### **Please do not:**

- Supply files that are optimized for screen use (e.g., GIF, BMP, PICT, WPG); these typically have a low number of pixels and limited set of colors;
- Supply files that are too low in resolution;
- Submit graphics that are disproportionately large for the content.

##### **Formats**

Regardless of the application used, when your electronic artwork is finalised, please "save as" or convert the images to one of the following formats (Note the resolution requirements for line drawings, halftones, and line/halftone combinations given below.):

EPS: Vector drawings. Embed the font or save the text as "graphics".

TIFF: Colour or greyscale photographs (halftones): always use a minimum of 300 dpi. For colour images always use RGB.

TIFF: Bitmapped line drawings: use a minimum of 1000 dpi.

TIFF: Combinations bitmapped line/half-tone (colour or greyscale): a minimum of 500 dpi is required.

DOC, XLS or PPT: If your electronic artwork is created in any of these Microsoft Office applications please supply "as is".

##### **Please do not:**

- Supply embedded graphics in your wordprocessor (spreadsheet, presentation) document;
- Supply files that are optimised for screen use (like GIF, BMP, PICT, WPG); the resolution is too low;
- Supply files that are too low in resolution;
- Submit graphics that are disproportionately large for the content.

##### *Color artwork*

Please make sure that artwork files are in an acceptable format (TIFF (or JPEG), EPS (or PDF), or MS Office files) and with the correct resolution. If, together with your accepted article, you submit usable color figures then Elsevier will ensure, at no additional charge, that these figures will appear in color online (e.g., ScienceDirect and other sites) regardless of whether or not these illustrations are reproduced in color in the printed version. **For color reproduction in print, you will receive**

**information regarding the costs from Elsevier after receipt of your accepted article.** Please indicate your preference for color: in print or online only. [Further information on the preparation of electronic artwork.](#)

#### *Figure captions*

Ensure that each illustration has a caption. Supply captions separately, not attached to the figure. A caption should comprise a brief title (**not** on the figure itself) and a description of the illustration. Keep text in the illustrations themselves to a minimum but explain all symbols and abbreviations used.

#### **Tables**

Please submit tables as editable text and not as images. Tables can be placed either next to the relevant text in the article, or on separate page(s) at the end. Number tables consecutively in accordance with their appearance in the text and place any table notes below the table body. Be sparing in the use of tables and ensure that the data presented in them do not duplicate results described elsewhere in the article. Please avoid using vertical rules and shading in table cells.

#### **References**

##### **Citation in text**

Please ensure that every reference cited in the text is also present in the reference list (and vice versa). Personal communications may be cited (with exact date) in the text but are not included in the reference list. Unpublished studies or papers may be cited but must include a date (year) and follow APA style. Citing reference as "in press" indicates that the work has been accepted for publication."

##### *Data references*

This journal encourages you to cite underlying or relevant datasets in your manuscript by citing them in your text and including a data reference in your Reference List. Data references should include the following elements: author name(s), dataset title, data repository, version (where available), year, and global persistent identifier. Add [dataset] immediately before the reference so we can properly identify it as a data reference. The [dataset] identifier will not appear in your published article.

##### *References in a special issue*

Please ensure that the words 'this issue' are added to any references in the list (and any citations in the text) to other articles in the same Special Issue.

##### *Reference management software*

Most Elsevier journals have their reference template available in many of the most popular reference management software products. These include all products that support [Citation Style Language styles](#), such as [Mendeley](#) and [Zotero](#), as well as [EndNote](#). Using the word processor plug-ins from these products, authors only need to select the appropriate journal template when preparing their article, after which citations and bibliographies will be automatically formatted in the journal's style. If no template is yet available for this journal, please follow the format of the sample references and citations as shown in this Guide.

Users of Mendeley Desktop can easily install the reference style for this journal by clicking the following link:

<http://open.mendeley.com/use-citation-style/body-image>

When preparing your manuscript, you will then be able to select this style using the Mendeley plug-ins for Microsoft Word or LibreOffice.

##### *Reference style*

**Text:** Citations in the text should follow the referencing style used by the American Psychological Association. You are referred to the Publication Manual of the American Psychological Association, Sixth Edition, ISBN 978-1-4338-0561-5, copies of which may be [ordered online](#) or APA Order Dept., P.O.B. 2710, Hyattsville, MD 20784, USA or APA, 3 Henrietta Street, London, WC3E 8LU, UK.

**List:** references should be arranged first alphabetically and then further sorted chronologically if necessary. More than one reference from the same author(s) in the same year must be identified by the letters 'a', 'b', 'c', etc., placed after the year of publication.

##### *Examples:*

Reference to a journal publication:

Van der Geer, J., Hanraads, J. A. J., & Lupton, R. A. (2010). The art of writing a scientific article. *Journal of Scientific Communications*, 163, 51–59.

Reference to a book:

Strunk, W., Jr., & White, E. B. (2000). *The elements of style*. (4th ed.). New York: Longman, (Chapter 4).



Reference to a chapter in an edited book:

Mettam, G. R., & Adams, L. B. (2009). How to prepare an electronic version of your article. In B. S. Jones, & R. Z. Smith (Eds.), *Introduction to the electronic age* (pp. 281–304). New York: E-Publishing Inc.

Reference to a website:

Cancer Research UK. Cancer statistics reports for the UK. (2003). <http://www.cancerresearchuk.org/aboutcancer/statistics/cancerstatsreport/> Accessed 13 March 2003.

Reference to a dataset:

[dataset] Oguro, M., Imahiro, S., Saito, S., Nakashizuka, T. (2015). *Mortality data for Japanese oak wilt disease and surrounding forest compositions*. Mendeley Data, v1. <https://doi.org/10.17632/xwj98nb39r.1>.

Reference to a conference paper or poster presentation:

Engle, E.K., Cash, T.F., & Jarry, J.L. (2009, November). The Body Image Behaviours Inventory-3: Development and validation of the Body Image Compulsive Actions and Body Image Avoidance Scales. Poster session presentation at the meeting of the Association for Behavioural and Cognitive Therapies, New York, NY.

### **Video**

Elsevier accepts video material and animation sequences to support and enhance your scientific research. Authors who have video or animation files that they wish to submit with their article are strongly encouraged to include links to these within the body of the article. This can be done in the same way as a figure or table by referring to the video or animation content and noting in the body text where it should be placed. All submitted files should be properly labeled so that they directly relate to the video file's content. In order to ensure that your video or animation material is directly usable, please provide the file in one of our recommended file formats with a preferred maximum size of 150 MB per file, 1 GB in total. Video and animation files supplied will be published online in the electronic version of your article in Elsevier Web products, including [ScienceDirect](#). Please supply 'stills' with your files: you can choose any frame from the video or animation or make a separate image. These will be used instead of standard icons and will personalize the link to your video data. For more detailed instructions please visit our [video instruction pages](#). Note: since video and animation cannot be embedded in the print version of the journal, please provide text for both the electronic and the print version for the portions of the article that refer to this content.

### **AudioSlides**

The journal encourages authors to create an AudioSlides presentation with their published article. AudioSlides are brief, webinar-style presentations that are shown next to the online article on ScienceDirect. This gives authors the opportunity to summarize their research in their own words and to help readers understand what the paper is about. [More information and examples are available](#). Authors of this journal will automatically receive an invitation e-mail to create an AudioSlides presentation after acceptance of their paper.

### **Data visualization**

Include interactive data visualizations in your publication and let your readers interact and engage more closely with your research. Follow the instructions [here](#) to find out about available data visualization options and how to include them with your article.

### **Supplementary material**

Supplementary material such as applications, images and sound clips, can be published with your article to enhance it. Submitted supplementary items are published exactly as they are received (Excel or PowerPoint files will appear as such online). Please submit your material together with the article and supply a concise, descriptive caption for each supplementary file. If you wish to make changes to supplementary material during any stage of the process, please make sure to provide an updated file. Do not annotate any corrections on a previous version. Please switch off the 'Track Changes' option in Microsoft Office files as these will appear in the published version.

### **Research data**

This journal encourages and enables you to share data that supports your research publication where appropriate, and enables you to interlink the data with your published articles. Research data refers to the results of observations or experimentation that validate research findings. To facilitate reproducibility and data reuse, this journal also encourages you to share your software, code, models, algorithms, protocols, methods and other useful materials related to the project.

Below are a number of ways in which you can associate data with your article or make a statement about the availability of your data when submitting your manuscript. If you are sharing data in one of these ways, you are encouraged to cite the data in your manuscript and reference list. Please refer to the "References" section for more information about data citation. For more information on depositing, sharing and using research data and other relevant research materials, visit the [research data](#) page.

#### *Data linking*

If you have made your research data available in a data repository, you can link your article directly to the dataset. Elsevier collaborates with a number of repositories to link articles on ScienceDirect with relevant repositories, giving readers access to underlying data that gives them a better understanding of the research described.

There are different ways to link your datasets to your article. When available, you can directly link your dataset to your article by providing the relevant information in the submission system. For more information, visit the [database linking page](#).

For [supported data repositories](#) a repository banner will automatically appear next to your published article on ScienceDirect.

In addition, you can link to relevant data or entities through identifiers within the text of your manuscript, using the following format: Database: xxxx (e.g., TAIR: AT1G01020; CCDC: 734053; PDB: 1XFN).

#### *Mendeley Data*

This journal supports Mendeley Data, enabling you to deposit any research data (including raw and processed data, video, code, software, algorithms, protocols, and methods) associated with your manuscript in a free-to-use, open access repository. During the submission process, after uploading your manuscript, you will have the opportunity to upload your relevant datasets directly to *Mendeley Data*. The datasets will be listed and directly accessible to readers next to your published article online.

For more information, visit the [Mendeley Data for journals page](#).

#### *Data statement*

To foster transparency, we encourage you to state the availability of your data in your submission. This may be a requirement of your funding body or institution. If your data is unavailable to access or unsuitable to post, you will have the opportunity to indicate why during the submission process, for example by stating that the research data is confidential. The statement will appear with your published article on ScienceDirect. For more information, visit the [Data Statement page](#).

### **AFTER ACCEPTANCE**

#### **Online proof correction**

Corresponding authors will receive an e-mail with a link to our online proofing system, allowing annotation and correction of proofs online. The environment is similar to MS Word: in addition to editing text, you can also comment on figures/tables and answer questions from the Copy Editor. Web-based proofing provides a faster and less error-prone process by allowing you to directly type your corrections, eliminating the potential introduction of errors.

If preferred, you can still choose to annotate and upload your edits on the PDF version. All instructions for proofing will be given in the e-mail we send to authors, including alternative methods to the online version and PDF.

We will do everything possible to get your article published quickly and accurately. Please use this proof only for checking the typesetting, editing, completeness and correctness of the text, tables and figures. Significant changes to the article as accepted for publication will only be considered at this stage with permission from the Editor. It is important to ensure that all corrections are sent back to us in one communication. Please check carefully before replying, as inclusion of any subsequent corrections cannot be guaranteed. Proofreading is solely your responsibility.

#### **Offprints**

The corresponding author will, at no cost, receive a customized [Share Link](#) providing 50 days free access to the final published version of the article on [ScienceDirect](#). The Share Link can be used for sharing the article via any communication channel, including email and social media. For an extra charge, paper offprints can be ordered via the offprint order form which is sent once the article is accepted for publication. Both corresponding and co-authors may order offprints at any time via

Elsevier's [Webshop](#). Corresponding authors who have published their article gold open access do not receive a Share Link as their final published version of the article is available open access on ScienceDirect and can be shared through the article DOI link.

#### **AUTHOR INQUIRIES**

Visit the [Elsevier Support Center](#) to find the answers you need. Here you will find everything from Frequently Asked Questions to ways to get in touch.

You can also [check the status of your submitted article](#) or find out [when your accepted article will be published](#).

© Copyright 2018 Elsevier | <https://www.elsevier.com>

## Appendix II: Copyright permission (The ARC Framework of Adjustment to Visible Difference) - Wiley

5/5/2018

RE: NON RIGHTSLINK - Permission to reproduce image - Kirsty Ryninks

RE: NON RIGHTSLINK - Permission to reproduce image

Wiley Global Permissions <permissions@wiley.com >

Wed 24/01/2018 11:33

To: Kirsty Ryninks <K.E.Ryninks@bath.ac.uk>;

Dear Kirsty,

Thank you for your email.

Permission is hereby granted for the use requested, subject to the usual acknowledgements (author, title of material, title of book/journal, ourselves as publisher). You should also duplicate the copyright notice that appears in the Wiley publication; this can be found on the copyright page if the material is a book or within the article if it is a journal.

Any third-party material is expressly excluded from this permission. If any of the material you wish to use appears within our work with credit to another source, authorisation from that source must be obtained.

This permission does not include the right to grant others permission to photocopy or otherwise reproduce this material except for accessible versions made by non-profit organisations serving the blind, visually impaired and other persons with print disabilities (VIPs).

Many thanks,

Orla Davies  
Rights Assistant  
John Wiley & Sons Ltd

**WILEY**

From: Kirsty Ryninks [mailto:K.E.Ryninks@bath.ac.uk]

Sent: 23 January 2018 17:10

To: Wiley Global Permissions <permissions@wiley.com>

Subject: NON RIGHTSLINK - Permission to reproduce image

From: Kirsty Ryninks

Sent: 23 January 2018 17:04

To: mailto:

Subject: Permission to reproduce image.

To whom it may concern,

I have been trying to request permission through your website to reproduce an image from one of your book titles but I am having difficulty locating the "Request Permissions" link as outlined in your guidance ([http://wileyrights.gms.sg/wiley\\_content.php](http://wileyrights.gms.sg/wiley_content.php)).

I would like to request permission to reproduce Figure A.11 on page 270 of your book titled "CBT for Appearance Anxiety: Psychosocial Interventions for Anxiety due to Visible Difference" by Alex Clarke et al.

The image will be included in my doctoral portfolio as part of my training at the University of Bath, as well as in a review article I have been writing with Dr Paul Salkovskis that we intend to submit for publication.

Would you be happy for us to include this image in our review article? And how can we go about obtaining this permission?

We look forward to hearing from you.

Kind Regards,  
Kirsty Ryninks

Clinical Psychologist in Training, University of Bath



### Appendix III: Copyright permission (The cognitive behavioural model of Body Dysmorphic Disorder) – David Veale

5/5/2018

Re: Permission to reproduce Figure - Kirsty Ryninks

Re: Permission to reproduce Figure

Kirsty Ryninks

Mon 22/01/2018 14:38

To: Veale, David <david.veale@kcl.ac.uk>;

Dear David,

Thank you for your email giving permission to reproduce and sending the book chapter.

I think we will reference the 2004 article as that is the image we have used in the development of our model. However, we would like to reference the 2017 chapter in the review itself. Thank you for bringing it to our attention.

Best Wishes,  
Kirsty

---

**From:** Veale, David <david.veale@kcl.ac.uk>

**Sent:** 21 January 2018 14:04

**To:** Kirsty Ryninks

**Subject:** Re: Permission to reproduce Figure

Yes of course although you may be interested in an updated book chapter published in 2017 see Figure 23.1  
Baldock E, Veale D. (2017). The self as an aesthetic object: a model of cognitive processing in Body Dysmorphic Disorder. Phillips, K.A. (editor). Oxford University Press. Chapter 23. pp. 299-310

David

---

**From:** Kirsty Ryninks <K.E.Ryninks@bath.ac.uk>

**Date:** Wednesday, 17 January 2018 at 14:52

**To:** David Veale <david.veale@kcl.ac.uk>

**Subject:** Permission to reproduce Figure

Dear David,

You may remember meeting with me, Cara Davis and Paul Salkovskis at Bath University to discuss the review article we have been working on looking at psychological adjustment to scarring following planned surgery.

We are planning to submit the article we have written to Body Image to be considered for publication and would like to request permission to reproduce Figure 1 (A cognitive behavioural model of BDD) from the following article:

Veale, D. (2004). Advances in a cognitive behavioural model of body dysmorphic disorder. *Body image*, 1(1), 113-125.

Please let us know if you are willing to grant this permission.

I look forward to hearing from you.

Kind Regards,  
Kirsty Ryninks

Clinical Psychologist in Training, University of Bath

## **Appendix IV: Instructions for Authors – British Journal of Clinical Psychology**

### ***Author Guidelines***

The British Journal of Clinical Psychology publishes original contributions to scientific knowledge in clinical psychology. This includes descriptive comparisons, as well as studies of the assessment, aetiology and treatment of people with a wide range of psychological problems in all age groups and settings. The level of analysis of studies ranges from biological influences on individual behaviour through to studies of psychological interventions and treatments on individuals, dyads, families and groups, to investigations of the relationships between explicitly social and psychological levels of analysis.

All papers published in The British Journal of Clinical Psychology are eligible for Panel A: Psychology, Psychiatry and Neuroscience in the Research Excellence Framework (REF).

The following types of paper are invited:

- Papers reporting original empirical investigations
- Theoretical papers, provided that these are sufficiently related to the empirical data
- Review articles which need not be exhaustive but which should give an interpretation of the state of the research in a given field and, where appropriate, identify its clinical implications
- Brief reports and comments

#### **1. Circulation**

The circulation of the Journal is worldwide. Papers are invited and encouraged from authors throughout the world.

#### **2. Length**

The word limit for papers submitted for consideration to BJCP is 5000 words and any papers that are over this word limit will be returned to the authors. The word limit does not include the abstract, reference list, figures, or tables. Appendices however are included in the word limit. The Editors retain discretion to publish papers beyond this length in cases where the clear and concise expression of the scientific content requires greater length. In such a case, the authors should contact the Editors before submission of the paper.

#### **3. Submission and reviewing**

All manuscripts must be submitted via [Editorial Manager](#). The Journal operates a policy of anonymous (double blind) peer review. We also operate a triage process in which submissions that are out of scope or otherwise inappropriate will be rejected by the editors without external peer review to avoid unnecessary delays. Before submitting, please read the [terms and conditions of submission](#) and the [declaration of competing interests](#). You may also like to use the [Submission Checklist](#) to help you prepare your paper.

#### **4. Manuscript requirements**

- Contributions must be typed in double spacing with wide margins. All sheets must be numbered.
- Manuscripts should be preceded by a title page which includes a full list of authors and their affiliations, as well as the corresponding author's contact details. You may like to

use [this](#) template. When entering the author names into Editorial Manager, the corresponding author will be asked to provide a CRediT contributor role to classify the role that each author played in creating the manuscript. Please see the [Project CRediT](#) website for a list of roles.

- The main document must be anonymous. Please do not mention the authors' names or affiliations (including in the Method section) and refer to any previous work in the third person.
- Tables should be typed in double spacing, each on a separate page with a self-explanatory title. Tables should be comprehensible without reference to the text. They should be placed at the end of the manuscript but they must be mentioned in the text.
- Figures can be included at the end of the document or attached as separate files, carefully labelled in initial capital/lower case lettering with symbols in a form consistent with text use. Unnecessary background patterns, lines and shading should be avoided. Captions should be listed on a separate sheet. The resolution of digital images must be at least 300 dpi. All figures must be mentioned in the text.
- All papers must include a structured abstract of up to 250 words under the headings: Objectives, Methods, Results, Conclusions. Articles which report original scientific research should also include a heading 'Design' before 'Methods'. The 'Methods' section for systematic reviews and theoretical papers should include, as a minimum, a description of the methods the author(s) used to access the literature they drew upon. That is, the abstract should summarize the databases that were consulted and the search terms that were used.
- All Articles must include Practitioner Points – these are 2–4 bullet points to detail the positive clinical implications of the work, with a further 2–4 bullet points outlining cautions or limitations of the study. They should be placed below the abstract, with the heading 'Practitioner Points'.
- For reference citations, please use APA style. Particular care should be taken to ensure that references are accurate and complete. Give all journal titles in full and provide DOI numbers where possible for journal articles.
- SI units must be used for all measurements, rounded off to practical values if appropriate, with the imperial equivalent in parentheses.
- In normal circumstances, effect size should be incorporated.
- Authors are requested to avoid the use of sexist language.
- Authors are responsible for acquiring written permission to publish lengthy quotations, illustrations, etc. for which they do not own copyright. For guidelines on editorial style, please consult the [APA Publication Manual](#) published by the American Psychological Association.

If you need more information about submitting your manuscript for publication, please email Vicki Pang, Editorial Assistant ([bjc@wiley.com](mailto:bjc@wiley.com)) or phone +44 (0) 1243 770 410 (ex 434 10).

## 5. Brief reports and comments

These allow publication of research studies and theoretical, critical or review comments with an essential contribution to make. They should be limited to 2000 words, including references. The abstract should not exceed 120 words and should be structured under these headings: Objective, Method, Results, Conclusions. There should be no more than one table

or figure, which should only be included if it conveys information more efficiently than the text. Title, author name and address are not included in the word limit.

## 6. Supporting Information

BJC is happy to accept articles with supporting information supplied for online only publication. This may include appendices, supplementary figures, sound files, videoclips etc. These will be posted on Wiley Online Library with the article. The print version will have a note indicating that extra material is available online. Please indicate clearly on submission which material is for online only publication. Please note that extra online only material is published as supplied by the author in the same file format and is not copyedited or typeset. Further information about this service can be found at <http://authorservices.wiley.com/bauthor/suppmat.asp>

## 7. Copyright and licenses

If your paper is accepted, the author identified as the formal corresponding author for the paper will receive an email prompting them to login into Author Services, where via the Wiley Author Licensing Service (WALS) they will be able to complete the license agreement on behalf of all authors on the paper.

### **For authors signing the copyright transfer agreement**

If the OnlineOpen option is not selected the corresponding author will be presented with the copyright transfer agreement (CTA) to sign. The terms and conditions of the CTA can be previewed in the samples associated with the [Copyright FAQs](#).

### **For authors choosing OnlineOpen**

If the OnlineOpen option is selected the corresponding author will have a choice of the following Creative Commons License Open Access Agreements (OAA):

- Creative Commons Attribution Non-Commercial License OAA
- Creative Commons Attribution Non-Commercial -NoDerivs License OAA

To preview the terms and conditions of these open access agreements please visit the [Copyright FAQs](#) and you may also like to visit the [Wiley Open Access Copyright and Licence](#) page.

If you select the OnlineOpen option and your research is funded by The Wellcome Trust and members of the Research Councils UK (RCUK) or the Austrian Science Fund (FWF) you will be given the opportunity to publish your article under a CC-BY license supporting you in complying with your Funder requirements. For more information on this policy and the Journal's compliant self-archiving policy please visit our [Funder Policy](#) page.

## 8. Colour illustrations

Colour illustrations can be accepted for publication online. These would be reproduced in greyscale in the print version. If authors would like these figures to be reproduced in colour in print at their expense they should request this by completing a Colour Work Agreement form upon acceptance of the paper. A copy of the Colour Work Agreement form can be downloaded [here](#).

## 9. Pre-submission English-language editing



Authors for whom English is a second language may choose to have their manuscript professionally edited before submission to improve the English. A list of independent suppliers of editing services can be found at [http://authorservices.wiley.com/bauthor/english\\_language.asp](http://authorservices.wiley.com/bauthor/english_language.asp). All services are paid for and arranged by the author, and use of one of these services does not guarantee acceptance or preference for publication.

#### 10. Author Services

Author Services enables authors to track their article – once it has been accepted – through the production process to publication online and in print. Authors can check the status of their articles online and choose to receive automated e-mails at key stages of production. The author will receive an e-mail with a unique link that enables them to register and have their article automatically added to the system. Please ensure that a complete e-mail address is provided when submitting the manuscript.

Visit <http://authorservices.wiley.com/bauthor/> for more details on online production tracking and for a wealth of resources including FAQs and tips on article preparation, submission and more.

#### 11. The Later Stages

The corresponding author will receive an email alert containing a link to a web site. A working e-mail address must therefore be provided for the corresponding author. The proof can be downloaded as a PDF (portable document format) file from this site. Acrobat Reader will be required in order to read this file. This software can be downloaded (free of charge) from the following web site: <http://www.adobe.com/products/acrobat/readstep2.html>.

This will enable the file to be opened, read on screen and annotated direct in the PDF. Corrections can also be supplied by hard copy if preferred. Further instructions will be sent with the proof. Excessive changes made by the author in the proofs, excluding typesetting errors, will be charged separately.

#### 12. Early View

British Journal of Clinical Psychology is covered by the Early View service on Wiley Online Library. Early View articles are complete full-text articles published online in advance of their publication in a printed issue. Articles are therefore available as soon as they are ready, rather than having to wait for the next scheduled print issue. Early View articles are complete and final. They have been fully reviewed, revised and edited for publication, and the authors' final corrections have been incorporated. Because they are in final form, no changes can be made after online publication. The nature of Early View articles means that they do not yet have volume, issue or page numbers, so they cannot be cited in the traditional way. They are cited using their Digital Object Identifier (DOI) with no volume and issue or pagination information. E.g., Jones, A.B. (2010). Human rights Issues. *Human Rights Journal*. Advance online publication. doi:10.1111/j.1467-9299.2010.00300.x

## Appendix V: University of Bath Service Improvement Project Ethical Approval

5/5/2018

RE: Ethics 16-148 - Kirsty Ryninks

RE: Ethics 16-148

psychology-ethics

Wed 15/06/2016 10:58

To: Kirsty Ryninks <K.E.Ryninks@bath.ac.uk>;

Dear Kirsty Ryninks

**Ethics Reference 16-148: Volunteers' experiences of helping hoarders' experiences of being helped**

Thank you for satisfactorily attending to those amendments. It appears your method of selected boxes in the final section confused the reviewers, but this is clarified now. I can now confirm that you have full ethical approval for your study.

Best wishes with your research,  
Dr Michael J Proulx  
Chair, Psychology Research Ethics Committee

5/5/2018

RE: Ethics Reference 16-148: Volunteers' experiences of helping hoarders' experiences of being helped - Kirsty Ryninks

RE: Ethics Reference 16-148: Volunteers' experiences of helping hoarders' experiences of being helped

psychology-ethics

Mon 24/10/2016 16:06

To: Kirsty Ryninks <K.E.Ryninks@bath.ac.uk>;

Dear Kirsty,

Thank you for sending the amended wording and explanation. I am happy to approve these amendments via Chair's Action.

Best of luck with your data collection,  
Dr. Nathalia Gjerose,  
Chair, Psychology Ethics Committee

## Appendix VI: Service Improvement Project Study Materials (Client and volunteer/case worker: Information Sheet, Consent Form, Demographic Questionnaires, Debrief Sheet)



Version 2: 13.10.16

Department of  
Psychology



Building 10 West 3.42  
Claverton Down  
Bath BA2 7AY

Email: k.e.ryninks@bath.ac.uk

### Making Space Project evaluation

#### *Client Information Sheet*

##### **Context**

WE Care and Repair have supported many people who have accumulated numerous possessions and this led them to develop the Making Space project in February 2015. Making Space uses a volunteer peer support model to work with clients in their own homes. The project has a practical focus on the physical environment to improve the safety and quality of life of adults who have accumulated numerous possessions in their homes. Volunteers work to motivate and support clients to 'make space' in their home.

##### **Why is this study being done?**

About 2-4% of the population experience difficulties associated with accumulating numerous possessions and it can cause many problems in people's lives, such as with their family, their ability to work and to maintain social contacts. People often find it difficult to obtain or seek help and WE Care and Repair appear to be an organisation that people have found they can engage with.

It is important that WE Care and Repair understand the experience of delivering and receiving their Making Space project from both staff/volunteers and clients. This will help the service to keep doing the things that it's doing well and change the things that it's not doing so well. Consequently, we are asking WE Care and Repair staff/volunteers and clients if they can spare a little of their time to tell us about their experiences.

Before you decide to take part, it is important for you to understand why the project is being done and what it will involve. Please take time to read the following information carefully and discuss it with others if you wish. Ask us if there is anything that is not clear or if you would like more information.

##### **Why have I been asked to take part?**

We are contacting you because we understand you have received support from WE Care and Repair. We would like you to join in an evaluation of the service that the Making Space project provides to clients through WE Care and Repair.

##### **Do I have to take part?**

No. Participation in this project is voluntary. If you do not wish to take part, this will not change the service that you are offered in any way. If you decide to take part and then later change your mind, either before you start the project, during it or afterwards, you can withdraw without giving your reasons, and, if you wish, your data will be destroyed up until the point of it being fully anonymised.

**What will I be asked to do if I take part?**

The project will involve completing two questionnaires about difficulties associated with accumulating numerous possessions (these should take about 10-15 minutes). You will then be asked to share your experience of receiving a service from WE Care and Repair in relation to accumulating numerous possessions in an interview (about 30-60 minutes) conducted by Kirsty Ryninks (Trainee Clinical Psychologist) at the St Philips office of WE Care and Repair, University of Bath or in your home, whichever is easiest for you. The interview will be recorded so that we can listen to what you have said at a later date and you would be welcome to receive a copy of the recording if you wanted one. The recording will be transcribed verbatim for analysis but identifiable details will be removed and all the information you provide will be anonymised.

**Will my experiences and reports be kept confidential?**

Yes. All information collected about you during the course of the project will be kept confidential and will conform to the Data Protection Act of 1998 with respect to data collection, storage and destruction. This means that all paper-based and electronic information will be locked and password protected with access restricted to project personnel and any information about you will have your name and address removed so that you cannot be identified from it.

We plan to report our findings in academic related journals and also present them to professionals at meetings, workshops and conferences. We would like to share the information we learn from this project so that other organisations can better support their clients who accumulate numerous possessions. This can make people a little apprehensive but we can assure you that you will not be identifiable in any reports or publications arising from the project.

**Are there any advantages/benefits from taking part?**

This study will not help you directly but the information collected from you and other participants should help to improve our understanding of the service provided by WE Care and Repair. Our aim from conducting this, and other studies like it, is to help develop and improve the services that WE Care and Repair offer to clients.

**Are there any disadvantages/risks from taking part?**

We consider there to be minimal disadvantages (e.g. the inconvenience of completing the questionnaires and/or being interviewed).

**What to do next if I'm interested or have some questions?**

If you would like to find out more about taking part or have any questions about anything that you have read, please return the attached opt-in form by email to [K.e.ryninks@bath.ac.uk](mailto:K.e.ryninks@bath.ac.uk) or by posting to Doctorate in Clinical Psychology, Building 10 West 3.42, Claverton Down, Bath, BA2 7AY. You may also find it helpful to discuss this project with WE Care and Repair and they can pass on your details for us to contact you directly.

Thank you very much for your time and your support in helping us to continue to improve the service that WE Care and Repair and the Making Space project provide.

**Making Space Project evaluation: understanding people with hoarding  
difficulties experiences of being helped and volunteers' experience**

***Making Space Volunteer/Case Worker Information Sheet***

**Context**

WE Care and Repair have supported numerous people with hoarding difficulties and this led them to develop the Making Space project in February 2015. Making Space uses a volunteer peer support model to work with adults that hoard in their own homes. The project has a practical focus on the physical environment to improve the safety and quality of life of adults who hoard. Volunteers work to motivate and support clients to de-clutter and improve their environment at home.

**Why is this study being done?**

About 2-4% of the population experience hoarding difficulties and it can cause many problems in people's lives, such as with their family, their ability to work and to maintain social contacts. People often find it difficult to seek help and WE Care and Repair appear to be an organisation that people have found they can engage with.

It is important that WE Care and Repair understand the experience of delivering and receiving their Making Space project from both staff/volunteers and clients. This will help the service to keep doing the things that it's doing well and change the things that it's not doing so well. Consequently, we are asking WE Care and Repair staff/volunteers and clients if they can spare a little of their time to tell us about their experiences.

Before you decide to take part, it is important for you to understand why the project is being done and what it will involve. Please take time to read the following information carefully and discuss it with others if you wish. Ask us if there is anything that is not clear or if you would like more information.

**Why have I been asked to take part?**

We are contacting you because we understand you are involved in working with hoarders through WE Care and Repair. We would like you to join in an evaluation of the service that the Making Space project provides to clients through WE Care and Repair.

**Do I have to take part?**

No. Participation in this project is voluntary. If you do not wish to take part, this will not change your role in the service in any way. If you decide to take part and then later change your mind, either



before you start the project, during it or afterwards, you can withdraw without giving your reasons, and, if you wish, your data will be destroyed up until the point of it being fully anonymised.

**What will I be asked to do if I take part?**

The project will involve completing two questionnaires about hoarding (these should take about 10-15 minutes). You will then be asked to share your experience of receiving a service from WE Care and Repair in relation to hoarding difficulties in an interview (about 30-60 minutes) conducted by Kirsty Ryninks (Trainee Clinical Psychologist) at the St Philips office of WE Care and Repair or in your home, whichever is easiest for you. The interview will be recorded so that we can listen to what you have said at a later date and you would be welcome to receive a copy of the recording if you wanted one. The recording will be transcribed verbatim for analysis but identifiable details will be removed and all the information you provide will be anonymised.

**Will my experiences and reports be kept confidential?**

Yes. All information collected about you during the course of the project will be kept confidential and will conform to the Data Protection Act of 1998 with respect to data collection, storage and destruction. This means that all paper-based and electronic information will be locked and password protected with access restricted to project personnel and any information about you will have your name and address removed so that you cannot be identified from it.

We plan to report our findings in academic/health related journals and also present them to health professionals at meetings, workshops and conferences. We would like to share the information we learn from this project so that other organisations can better support their clients who hoard. This can make people a little apprehensive but we can assure you that people will not be identifiable in any reports or publications arising from the project.

**Are there any advantages/benefits from taking part?**

This study will not help you directly but the information collected from you and other participants should help to improve our understanding of the service provided by WE Care and Repair. Our aim from conducting this, and other studies like it, is to help develop and improve the hoarding related services that WE Care and Repair offer to clients.

**Are there any disadvantages/risks from taking part?**

We consider there to be minimal disadvantages (e.g. the inconvenience of completing the questionnaires and/or being interviewed).

**What to do next if I'm interested or have some questions?**

If you would like to find out more about taking part or have any questions about anything that you have read, please return the attached opt-in form by email to [K.e.ryninks@bath.ac.uk](mailto:K.e.ryninks@bath.ac.uk) or by posting to Doctorate in Clinical Psychology, Building 10 West 3.42, Claverton Down, Bath, BA2 7AY. You may also find it helpful to discuss this project with WE Care and Repair and they can pass on your details for us to contact you directly.

Thank you very much for your time and your support in helping us to continue to improve the service that WE Care and Repair and the Making Space project provide.



Version 2: 13.10.16

Department of  
Psychology



UNIVERSITY OF  
**BATH**

Building 10 West 3.42  
Claverton Down  
Bath BA2 7AY

Email: [k.e.ryninks@bath.ac.uk](mailto:k.e.ryninks@bath.ac.uk)

## **Making Space Project evaluation**

### ***Opt-in Form***

WE Care and Repair is working in partnership with the University of Bath. We would like you to join in an evaluation of the service that the Making Space project provides through WE Care and Repair.

If you are interested in taking part in this project and participating in a brief interview about your experience, please tick the box and fill in your contact details.

☐

I would be happy to be contacted by Kirsty Ryninks (Trainee Clinical Psychologist) using the information I have provided below.

**Name:** \_\_\_\_\_

**Email:** \_\_\_\_\_

**Telephone:** \_\_\_\_\_

**Signature:** \_\_\_\_\_

Please return this form by email to [k.e.ryninks@bath.ac.uk](mailto:k.e.ryninks@bath.ac.uk) or by posting to Doctorate in Clinical Psychology, Building 10 West 3.42, Claverton Down, Bath, BA2 7AY.

Thank you very much for your time and your support in helping us to continue to improve the service that WE Care and Repair and the Making Space project provide.

## Making Space Project evaluation:

### Consent Form

	<b>Please initial all boxes</b>
1. I confirm that I have read and understood the information sheet dated 13.10.16 (version 2) for the above project. I have had the opportunity to consider the information, ask questions, and have had these answered satisfactorily.	<input type="checkbox"/>
2. I understand that my participation is voluntary, that I am free to withdraw at any time without having to give a reason, and that my decision will not affect my employment/care/legal rights.	<input type="checkbox"/>
3. I understand that any information about me will be handled in confidence. Any analyses and reports or publications (including short quotations in the media) from the data will be summary data only and will not identify me.	<input type="checkbox"/>
4. I give consent for my interview to be taped for the purposes of later transcription and for the researchers to ensure the quality of the interviewing process.	<input type="checkbox"/>
5. I agree to take part in the project.	<input type="checkbox"/>

\_\_\_\_\_  
Name of Participant      Date      Signature

\_\_\_\_\_  
Name of person      Date      Signature  
taking consent

Thank you very much for your time and your support in helping us to continue to improve the service that WE Care and Repair and the Making Space project provide.



**Making Space Project evaluation:**

***Demographic Questionnaire – Client***

To help us understand the information from the interview, please tell us a little bit about you.

Please complete/tick the relevant boxes below.

<b>Age</b>	_____ years old
<b>Gender</b>	<input type="checkbox"/> Male <input type="checkbox"/> Female
<b>Marital Status</b>	<input type="checkbox"/> Single <input type="checkbox"/> In Relationship <input type="checkbox"/> Married <input type="checkbox"/> Separated <input type="checkbox"/> Divorced <input type="checkbox"/> Widowed
<b>Employment Status</b>	<input type="checkbox"/> Full-time <input type="checkbox"/> Part-time <input type="checkbox"/> Retired <input type="checkbox"/> Home maker <input type="checkbox"/> Looking for work/unemployed <input type="checkbox"/> Other (specify): _____
<b>What is your ethnic group?</b>	<input type="checkbox"/> White UK <input type="checkbox"/> White Irish <input type="checkbox"/> Black Caribbean <input type="checkbox"/> Black African <input type="checkbox"/> Chinese <input type="checkbox"/> Bangladeshi <input type="checkbox"/> Indian <input type="checkbox"/> Pakistani <input type="checkbox"/> Mixed/Multiple Ethnic Background (specify): _____ <input type="checkbox"/> Other (specify): _____

<b>What is your highest level of Education?</b>	<input type="checkbox"/> GCSE or equivalent <input type="checkbox"/> A-level or equivalent <input type="checkbox"/> Bachelors degree <input type="checkbox"/> Masters degree <input type="checkbox"/> Other (specify): _____
<b>Does anyone else live with you at home?</b>	<input type="checkbox"/> Yes <input type="checkbox"/> No If Yes, please give details:
<b>Do you think you have/have had a problem accumulating numerous possessions?</b>	<input type="checkbox"/> Yes <input type="checkbox"/> No If Yes, when do you think this became a problem?  <input type="checkbox"/> Childhood <input type="checkbox"/> Adolescence <input type="checkbox"/> In my 20s <input type="checkbox"/> In my 30s <input type="checkbox"/> In my 40s <input type="checkbox"/> In my 50s <input type="checkbox"/> In my 60s <input type="checkbox"/> In my 70s <input type="checkbox"/> Other (specify): _____
<b>Have you ever received help from a health professional for problems relating to accumulating numerous possessions?</b>	<input type="checkbox"/> Yes <input type="checkbox"/> No  If Yes, how many times have you received help and when was this?    What kind of support did you receive and for how long?    Overall, how helpful was this support on a scale of 1 to 10, 1 being "not helpful at all" and 10 being "extremely helpful"?

<p><b>Are you currently taking any medication in relation to difficulties accumulating numerous possessions?</b></p>	<p><input type="checkbox"/> Yes <input type="checkbox"/> No</p> <p>(Please Specify if Yes):</p>
<p><b>How many visits did you have with a Making Space volunteer?</b></p>	
<p><b>How long, on average, did the visits last?</b></p>	
<p><b>Over approximately how many months did your visits last?</b></p>	
<p><b>Overall, how helpful were the visits with the Making Space volunteer on a scale of 1 to 10, 1 being "not helpful at all" and 10 being "extremely helpful"?</b></p>	

Thank you very much for your time and your support in helping us to continue to improve the service that WE Care and Repair and the Making Space project provide.

**Making Space Project evaluation: understanding people with hoarding  
difficulties experiences of being helped and volunteers' experience:**

***Demographic Questionnaire – Making Space Volunteer/Case Worker***

To help us understand the information from the interview we need to know a little bit about you.

Please complete/tick the relevant boxes below.

<b>Age</b>	_____ years old
<b>Gender</b>	<input type="checkbox"/> Male <input type="checkbox"/> Female
<b>Employment Status</b>	<input type="checkbox"/> Full-time <input type="checkbox"/> Part-time <input type="checkbox"/> Retired <input type="checkbox"/> Home maker <input type="checkbox"/> Looking for work/unemployed <input type="checkbox"/> Other (specify): _____
<b>What is/was your occupation?</b>	
<b>What is your ethnic group?</b>	<input type="checkbox"/> White UK <input type="checkbox"/> White Irish <input type="checkbox"/> Black Caribbean <input type="checkbox"/> Black African <input type="checkbox"/> Chinese <input type="checkbox"/> Bangladeshi <input type="checkbox"/> Indian <input type="checkbox"/> Pakistani <input type="checkbox"/> Mixed/Multiple Ethnic Background (specify): _____ <input type="checkbox"/> Other (specify): _____

<b>What is your highest level of Education?</b>	<input type="checkbox"/> GCSE or equivalent <input type="checkbox"/> A-level or equivalent <input type="checkbox"/> Bachelors degree <input type="checkbox"/> Masters degree <input type="checkbox"/> Other (specify): _____
<b>Did you have any experience of working with people with hoarding difficulties prior to becoming a case worker/volunteer?</b>	
<b>How long have you been a case worker/volunteer with the Making Space project?</b>	
<b>Have you undertaken any training that has helped you to work with clients in the Making Space project?</b>	<input type="checkbox"/> Yes <input type="checkbox"/> No If Yes, please give details:
<b>Approximately how many clients have you worked with?</b>	

Thank you very much for your time and your support in helping us to continue to improve the service that WE Care and Repair and the Making Space project provide.

## Making Space Project evaluation

### Debrief Sheet

Thank you for taking part in this project. WE Care and Repair have supported many people who have accumulated numerous possessions and want to understand the experience of delivering and receiving their Making Space project from both the perspective of their staff/volunteers and clients. Your support in this project will help the service to keep doing the things that it's doing well and change the things that it's not.

If you would like to receive a written summary of the findings of the project when that becomes available, please either email [K.E.Ryninks@bath.ac.uk](mailto:k.e.ryninks@bath.ac.uk) or ask WE Care and Repair to request this.

We have prepared a list of resources for participants who have taken part in the project so that you are aware of what options are available if you feel you would like some additional support. Please note that we do not expect you to be looking for support but we wanted to keep this list as comprehensive as possible. The list is not exhaustive however and we cannot specifically recommend each book/website/service or guarantee the quality of each.

#### Books and Websites

*Overcoming Compulsive Hoarding: Why you Save & How you Can Stop*, by Fugen Neziroglu, Jerome Bubrick and Jose Yaryura-Tobias.

[www.helpforhoarders.co.uk](http://www.helpforhoarders.co.uk)

Help for Hoarders - this informative site was developed to provide information, support and advice and create awareness about this secretive condition, for hoarders and their loved ones. The website contains a resources page with additional resources to those listed here.

[www.hoardinguk.org](http://www.hoardinguk.org)

Hoarding UK offer phone, email and advocacy support free of charge.

Thank you very much for your time and your support in helping us to continue to improve the service that WE Care and Repair and the Making Space project provide.

## **Appendix VII: Service Improvement Project Outcomes**

As the charity is currently undergoing a reorganisation and retendering process, providing formal feedback on this project has been difficult due to other more pressing priorities for the charity. The charity has received a copy of the project report and feedback has been given informally by email and telephone calls.

The charity has been contacted regarding presenting the findings to volunteers within the Making Space project and the wider organisation and this formal feedback will be given at their volunteer meeting on 12th September 2018.

The informal feedback that has been given to the charity to date has resulted in the following:

- Quotations from the interviews with clients and volunteers have been used in promotional material about the Making Space project and to support applications for additional funding for the project.
- The training package for volunteers is under review and the charity plans to include additional information on manual handling/hygiene in the future. The charity are also exploring the possibility of further training on the psychological aspects of working with compulsive hoarders.
- A peer support group for volunteers has been established and the feedback to date has been positive, with volunteers feeling more supported in their work with hoarders.
- The data collected from the CIRS and SI-R has been of interest to the charity and a consultancy project is currently exploring how to incorporate outcome measures into the project in the future.

## Appendix VIII: Instructions for Authors –BMC Pregnancy and Childbirth

### Research article

#### **Criteria**

Research articles should report on original primary research, but may report on systematic reviews of published research provided they adhere to the appropriate reporting guidelines which are detailed in our [editorial policies](#). Please note that non-commissioned pooled analyses of selected published research will not be considered.

*BMC Pregnancy and Childbirth* strongly encourages that all datasets on which the conclusions of the paper rely should be available to readers. We encourage authors to ensure that their datasets are either deposited in publicly available repositories (where available and appropriate) or presented in the main manuscript or additional supporting files whenever possible. Please see Springer Nature's [information on recommended repositories](#). Where a widely established research community expectation for data archiving in public repositories exists, submission to a community-endorsed, public repository is mandatory. A list of data where deposition is required, with the appropriate repositories, can be found on the [Editorial Policies Page](#).

Authors who need help depositing and curating data may wish to consider uploading their data to [Springer Nature's Research Data Support](#) or contacting our [Research Data Support Helpdesk](#). Springer Nature's Research Data Support provides data deposition and curation to help authors follow good practice in sharing and archiving of research data, and can be accessed [via an online form](#). The services provide secure and private submission of data files, which are curated and managed by the Springer Nature Research Data team for public release, in agreement with the submitting author. These services are provided in partnership with figshare. Checks are carried out as part of a submission screening process to ensure that researchers who should use a specific community-endorsed repository are advised of the best option for sharing and archiving their data. Use of Research Data Support is optional and does not imply or guarantee that a manuscript will be accepted.

#### **Preparing your manuscript**

The information below details the section headings that you should include in your manuscript and what information should be within each section.

Please note that your manuscript must include a 'Declarations' section including all of the subheadings (please see below for more information).



## **Title page**

The title page should:

- present a title that includes, if appropriate, the study design e.g.:
  - "A versus B in the treatment of C: a randomized controlled trial", "X is a risk factor for Y: a case control study", "What is the impact of factor X on subject Y: A systematic review"
  - or for non-clinical or non-research studies a description of what the article reports
- list the full names, institutional addresses and email addresses for all authors
  - if a collaboration group should be listed as an author, please list the Group name as an author. If you would like the names of the individual members of the Group to be searchable through their individual PubMed records, please include this information in the “Acknowledgements” section in accordance with the instructions below
- indicate the corresponding author

## **Abstract**

The Abstract should not exceed 350 words. Please minimize the use of abbreviations and do not cite references in the abstract. Reports of randomized controlled trials should follow the [CONSORT](#) extension for abstracts. The abstract must include the following separate sections:

- **BACKGROUND:** the context and purpose of the study
- **METHODS:** how the study was performed and statistical tests used
- **RESULTS:** the main findings
- **CONCLUSIONS:** brief summary and potential implications
- **TRIAL REGISTRATION:** If your article reports the results of a health care intervention on human participants, it must be registered in an appropriate registry and the registration number and date of registration should be in stated in this section. If it was not registered prospectively (before enrollment of the first participant), you should include the words 'retrospectively registered'. See our [editorial policies](#) for more information on trial registration

## **Keywords**

Three to ten keywords representing the main content of the article.

## **Background**

The Background section should explain the background to the study, its aims, a summary of the existing literature and why this study was necessary or its contribution to the field.

## **Methods**

The methods section should include:

- the aim, design and setting of the study
- the characteristics of participants or description of materials
- a clear description of all processes, interventions and comparisons. Generic drug names should generally be used. When proprietary brands are used in research, include the brand names in parentheses
- the type of statistical analysis used, including a power calculation if appropriate

## **Results**

This should include the findings of the study including, if appropriate, results of statistical analysis which must be included either in the text or as tables and figures.

## **Discussion**

This section should discuss the implications of the findings in context of existing research and highlight limitations of the study.

## **Conclusions**

This should state clearly the main conclusions and provide an explanation of the importance and relevance of the study reported.

## **List of abbreviations**

If abbreviations are used in the text they should be defined in the text at first use, and a list of abbreviations should be provided.

## **Declarations**

All manuscripts must contain the following sections under the heading 'Declarations':

- Ethics approval and consent to participate
- Consent for publication
- Availability of data and material
- Competing interests
- Funding
- Authors' contributions
- Acknowledgements
- Authors' information (optional)

Please see below for details on the information to be included in these sections.

If any of the sections are not relevant to your manuscript, please include the heading and write 'Not applicable' for that section.

### **Ethics approval and consent to participate**

Manuscripts reporting studies involving human participants, human data or human tissue must:

- include a statement on ethics approval and consent (even where the need for approval was waived)
- include the name of the ethics committee that approved the study and the committee's reference number if appropriate

Studies involving animals must include a statement on ethics approval.

See our [editorial policies](#) for more information.

If your manuscript does not report on or involve the use of any animal or human data or tissue, please state “Not applicable” in this section.

### **Consent for publication**

If your manuscript contains any individual person's data in any form (including individual details, images or videos), consent for publication must be obtained from that person, or in the case of children, their parent or legal guardian. All presentations of case reports must have consent for publication.

You can use your institutional consent form or our [consent form](#) if you prefer. You should not send the form to us on submission, but we may request to see a copy at any stage (including after publication).

See our [editorial policies](#) for more information on consent for publication.

If your manuscript does not contain data from any individual person, please state “Not applicable” in this section.

### **Availability of data and materials**

All manuscripts must include an ‘Availability of data and materials’ statement. Data availability statements should include information on where data supporting the results reported in the article can be found including, where applicable, hyperlinks to publicly archived datasets analysed or generated during the study. By data we mean the minimal dataset that would be necessary to interpret, replicate and build upon the findings reported in the article. We recognise it is not always possible to share research data publicly, for instance when individual privacy could be compromised, and in such instances data availability should still be stated in the manuscript along with any conditions for access.

Data availability statements can take one of the following forms (or a combination of more than one if required for multiple datasets):

- The datasets generated and/or analysed during the current study are available in the [NAME] repository, [PERSISTENT WEB LINK TO DATASETS]
- The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.
- All data generated or analysed during this study are included in this published article [and its supplementary information files].
- The datasets generated and/or analysed during the current study are not publicly available due [REASON WHY DATA ARE NOT PUBLIC] but are available from the corresponding author on reasonable request.
- Data sharing is not applicable to this article as no datasets were generated or analysed during the current study.
- The data that support the findings of this study are available from [third party name] but restrictions apply to the availability of these data, which were used under license for the current study, and so are not publicly available. Data are however available from the authors upon reasonable request and with permission of [third party name].
- Not applicable. If your manuscript does not contain any data, please state 'Not applicable' in this section.

More examples of template data availability statements, which include examples of openly available and restricted access datasets, are available [here](#).

BioMed Central also requires that authors cite any publicly available data on which the conclusions of the paper rely in the manuscript. Data citations should include a persistent identifier (such as a DOI) and should ideally be included in the reference list. Citations of datasets, when they appear in the reference list, should include the minimum information recommended by DataCite and follow journal style. Dataset identifiers including DOIs should be expressed as full URLs. For example:

Hao Z, AghaKouchak A, Nakhjiri N, Farahmand A. Global integrated drought monitoring and prediction system (GIDMaPS) data sets. figshare.  
2014. <http://dx.doi.org/10.6084/m9.figshare.853801>

With the corresponding text in the Availability of data and materials statement:

The datasets generated during and/or analysed during the current study are available in the [NAME] repository, [PERSISTENT WEB LINK TO DATASETS].<sup>[Reference number]</sup>

### **Competing interests**

All financial and non-financial competing interests must be declared in this section.

See our [editorial policies](#) for a full explanation of competing interests. If you are unsure whether you or any of your co-authors have a competing interest please contact the editorial office.

Please use the authors initials to refer to each author's competing interests in this section.

If you do not have any competing interests, please state "The authors declare that they have no competing interests" in this section.

## **Funding**

All sources of funding for the research reported should be declared. The role of the funding body in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript should be declared.

## **Authors' contributions**

The individual contributions of authors to the manuscript should be specified in this section. Guidance and criteria for authorship can be found in our [editorial policies](#).

Please use initials to refer to each author's contribution in this section, for example: "FC analyzed and interpreted the patient data regarding the hematological disease and the transplant. RH performed the histological examination of the kidney, and was a major contributor in writing the manuscript. All authors read and approved the final manuscript."

## **Acknowledgements**

Please acknowledge anyone who contributed towards the article who does not meet the criteria for authorship including anyone who provided professional writing services or materials.

Authors should obtain permission to acknowledge from all those mentioned in the Acknowledgements section.

See our [editorial policies](#) for a full explanation of acknowledgements and authorship criteria.

If you do not have anyone to acknowledge, please write "Not applicable" in this section.

Group authorship (for manuscripts involving a collaboration group): if you would like the names of the individual members of a collaboration Group to be searchable through their individual PubMed records, please ensure that the title of the collaboration Group is included on the title page and in the submission system and also include collaborating author names as the last paragraph of the "Acknowledgements" section. Please add authors in the format First Name, Middle initial(s) (optional), Last Name. You can add institution or

country information for each author if you wish, but this should be consistent across all authors.

Please note that individual names may not be present in the PubMed record at the time a published article is initially included in PubMed as it takes PubMed additional time to code this information.

### **Authors' information**

This section is optional.

You may choose to use this section to include any relevant information about the author(s) that may aid the reader's interpretation of the article, and understand the standpoint of the author(s). This may include details about the authors' qualifications, current positions they hold at institutions or societies, or any other relevant background information. Please refer to authors using their initials. Note this section should not be used to describe any competing interests.

### **Endnotes**

Endnotes should be designated within the text using a superscript lowercase letter and all notes (along with their corresponding letter) should be included in the Endnotes section. Please format this section in a paragraph rather than a list.

### **References**

All references, including URLs, must be numbered consecutively, in square brackets, in the order in which they are cited in the text, followed by any in tables or legends. The reference numbers must be finalized and the reference list fully formatted before submission.

## Appendix IX: Permission to use questionnaires from Posttraumatic Growth Research Centre

5/5/2018

Re: Request of copy of PTGI for Doctorate in Clinical Psyc... - Kirsty Ryninks

Re: Request of copy of PTGI for Doctorate in Clinical Psychology Research project

plewis19@uncc.edu on behalf of Posttraumatic Growth <posttraumaticgrowth@uncc.edu>

Mon 27/03/2017 20:45

To: Kirsty Ryninks <K.E.Ryninks@bath.ac.uk>;

Hello Ms. Ryninks,

Thank you for checking. You have our permission to use all of our scales in your research.

Warm regards,

Posttraumatic Growth Research Center

UNC Charlotte

Department of Psychology

9201 University City Blvd

Charlotte, NC 28223-0001 USA

Lawrence G. Calhoun ([lcalhoun@uncc.edu](mailto:lcalhoun@uncc.edu))

Richard G. Tedeschi ([rtedesch@uncc.edu](mailto:rtedesch@uncc.edu))

Arnie Cann ([acann@uncc.edu](mailto:acann@uncc.edu))

[www.ptg.uncc.edu](http://www.ptg.uncc.edu)

<http://www.routledge/mentalhealth.com/books/details/9780415645300/>

## Appendix X: University of Bath Main Research Project Ethical Approval

5/5/2018 Ethics 17-044 - Kirsty Ryninks

Ethics 17-044

Nathalia Gjersoe <N.Gjersoe@imaps.bath.ac.uk> on behalf of psychology-ethics <psychology-ethics@bath.ac.uk>  
Fri 10/03/2017 13:12  
To: Kirsty Ryninks <K.E.Ryninks@bath.ac.uk>;

Dear Kirst Ryninks,

Reference Number **16-044: Emotional Reactions to Loss in Pregnancy**

The ethics committee have considered your ethics proposal for the study above and have given it full ethical approval.

Best wishes with your research.

Dr Nathalia Gjersoe  
Chair, Psychology Research Ethics Committee

5/6/2018 Re: Ethics 17-044 - Kirsty Ryninks

Re: Ethics 17-044

Kirsty Ryninks  
Wed 15/11/2017 09:29  
To: Nathalia Gjersoe <N.Gjersoe@bath.ac.uk>;

Dear Nathalia,

Thank you for passing on the amendment instructions and apologies for contacting you directly. I will email the Psychology ethics address for any future changes and write an outline of the changes I am proposing.

The online survey has been amended to reflect the changes outlined in the information sheet, diversion page and demographic questionnaire, although it isn't possible to see the changes in the PDF I sent you. We have an online and paper version of our project so the documents are the same.

Thank you for approving the changes.

Best Wishes,  
Kirsty

---

**From:** Nathalia Gjersoe  
**Sent:** 15 November 2017 09:23  
**To:** Kirsty Ryninks  
**Subject:** RE: Ethics 17-044

Dear Kirsty,

Sorry for the delay. In future please follow the instructions for submitting amendments on the psychology moodle page. These should be submitted to [psychology-ethics@bath.ac.uk](mailto:psychology-ethics@bath.ac.uk), not my personal address (primarily because I keep on top of that account better so it will get seen to faster) and the amendments should be briefly explained in the covering email to ensure I am aware of all of the things you would like to draw my attention to.

I can't see any amendments to the online survey – is that correct? I am happy to confirm that you have full ethical approval for the amendments noted on the Information sheet, Diversion page, Demographic Questionnaire and Study Advert.

Best wishes,  
Dr. Nathalia Gjersoe  
Chair, Psychology Ethics Committee



## Appendix XI: Main Research Project Study Materials (Information Sheet, Consent Form, Demographic Questionnaire, Study Questionnaires, Debrief Sheet)

Version 3: 07.11.17

Department of  
Psychology



### EMOTIONAL REACTIONS TO LOSS IN PREGNANCY

#### *Information Sheet*

##### **Context**

We would like to invite you to take part in our research project exploring emotional reactions to loss in pregnancy. Before you decide whether or not you would like to take part, it is important to understand why the research is being done and what it would involve. Please take time to read the following information carefully and discuss it with friends and family if you wish. If you have any questions about this project, please contact the research team on [erlp@bath.ac.uk](mailto:erlp@bath.ac.uk)

##### **Why is this study being done?**

Miscarriage and stillbirth are unique and painful experiences and each person will experience different emotional reactions. We are interested in understanding more about these emotional reactions. We hope that the outcomes of this project will improve our understanding and enable us to improve the help and support offered to women and their families following early miscarriage and stillbirth.

##### **Who can take part in this project?**

We know that all parents' experiences of the death of their baby or babies are important to understand and explore. But to make sure we are asking good research questions and that we can make good sense of our findings, we have to be very specific about who takes part in this project. We are therefore focusing on certain groups of women and certain types of loss in this project. As we are following on from some earlier research, we are using the same timeframe that was used in the original work.

We are particularly interested in understanding the emotional reactions of the following groups of women in the UK and who are currently over the age of 18 years old:

- Women who experienced a **stillbirth (at 24 weeks of pregnancy or later)** 2 to 6 years ago
- Women who experienced an **early miscarriage (up to 12 weeks of pregnancy)** 2 to 6 years ago

Unfortunately, we are unable to invite the following groups to take part in this project at this time:

- Women who are currently under the age of 18 years
- Women who cannot read English because we are unable to translate the questionnaires
- Women who experienced a late miscarriage (between the 13th and 23rd week of pregnancy) 2 to 6 years ago
- Women who lost their baby more recently than 2 years or more than 6 years ago\*\* (see below)

\*\*Women who have had a pregnancy loss (early miscarriage or stillbirth) 2 to 6 years ago **and** a more recent loss **are able** to take part (as we will ask about the loss that affected you most in the last 24 months to 6 years).

[N. B. Women who have had a more recent loss but have not lost a baby in the 2-6 year time frame are not eligible for this particular study, but we hope we can do more research in this area in the future].

**What will I be asked to do if I take part?**

The project will involve completing a number of questionnaires online about yourself, your pregnancy loss and your emotional wellbeing. If you would prefer to complete the questionnaires by hand, please email the research team at [erlp@bath.ac.uk](mailto:erlp@bath.ac.uk) with your address and a copy of the survey questions will be sent to you in the post to complete anonymously and return in a stamped addressed envelope. Completing the survey is expected to take 30 minutes.

If you complete the survey online, we suggest that you use a device that you can easily type on and complete the survey at a time when you can be quiet and will not be disturbed. You will have the option to have a break and come back to the online survey by clicking on the "finish later" option at the bottom of each page. This will enable you to either bookmark the survey or to have the details emailed to you. All responses will be anonymous.

At the end of the study, we will ask some brief questions about your experience of taking part in the research. It is important to remember that there are no right or wrong answers to the questions as each individual's experience will be different.

**Do I have to take part?**

Participation in this project is voluntary. It is up to you to decide whether you wish to take part in the project or not. You can contact the research team by emailing [erlp@bath.ac.uk](mailto:erlp@bath.ac.uk) if you would like to speak further before deciding whether or not you would like to take part. We aim to respond to all emails within a week.

**Are there any advantages/benefits from taking part?**

We cannot promise that this project will help you directly but we hope that the information we collect will improve our understanding of emotional reactions following early miscarriage and stillbirth and improve the help and support that is offered to women and their families in the future. Previous research has shown that some people have found it helpful to take part in studies about their experiences and felt as though they were helping others by sharing their experiences. On completion of the project, a summary of our findings will be made available to participants who wish to receive them. You can request this by emailing [erlp@bath.ac.uk](mailto:erlp@bath.ac.uk).

**Are there any disadvantages/risks from taking part?**

Completing our survey will require you to think about thoughts and feelings that some people might find challenging. It is possible that you may find answering questions about your loss upsetting. This is a normal reaction when thinking about difficult experiences and such feelings will normally subside within a short time. It is important for you to understand that you are not required to answer questions or share anything that you do not want to.

If you are worried about your wellbeing, please make an appointment with your General Practitioner (GP). You could also contact other services that offer support and a list of helpful addresses and resources is available at the end of this information sheet. All your responses will be anonymous and so it will not be possible for the research team to identify who you are and whether your answers suggest you may be feeling anxious or depressed. If you have any concerns about your participation in the research, please contact a member of the research team by emailing [erlp@bath.ac.uk](mailto:erlp@bath.ac.uk). If you wish, we can arrange for someone to call you to discuss your concerns within three working days.

**Will my experiences be kept confidential?**

Yes. All information collected about you during the course of the project will be kept confidential and will conform to the Data Protection Act of 1998 with respect to data collection, storage and destruction. This means that all information will be locked and password protected with access restricted to project personnel.

You will not be asked to provide your name or contact details if you take part in the study so your data (whether completed online or by post) will be anonymous. If you email the research team to ask for information about the study, only the research team will know your contact details and this will be kept on a password-protected memory stick.

**Can I withdraw from the study?**

You are free to stop responding to the questions and your data will not be submitted *up until you have completed the survey or returned the survey in the post*. Once you have finished the survey online or anonymously posted your questionnaires to the research team, your data will be anonymous and it will not be possible to identify and withdraw your data.

**What will happen to the results of the project?**

This project is intended to form part of a doctoral thesis and we plan to report our findings in scientific journals and also present them to health professionals at meetings, workshops and conferences. This can make people a little apprehensive but we can assure you that the survey is anonymous and individual participants will not be identifiable in any written reports or publications arising from the project. We would like to share the information we learn from this project so that other organisations can better support women who have experienced early miscarriage and stillbirth.

**PART 2: What if there is a problem?**

Please read the following information where we provide detailed information about who to go to if you have any concerns or wish to complain about any aspect of the way you have been approached or treated as part of this project.

**Who is organising and funding this research? Who has reviewed this project?**

The project is being organised and funded by the University of Bath as part of the Doctorate in Clinical Psychology for Kirsty Ryninks. This research has received ethical approval from the University of Bath Psychology Ethics Committee under reference (17-044).

**What do I do if I have concerns about the project?**

Every care will be taken to ensure your safety during the course of the study. If you have any concerns about the project, please contact a member of the research team below by emailing [erlp@bath.ac.uk](mailto:erlp@bath.ac.uk):

CHIEF INVESTIGATOR – Kirsty Ryninks (Clinical Psychologist in Training, University of Bath)

ACADEMIC SUPERVISORS – Dr Megan Wilkinson-Tough (University of Bath), Dr Antje Horsch (University of Lausanne, Switzerland)

If you wish to complain formally, you can contact Professor Jonathan Knight (Pro-Vice Chancellor for Research) c/o Dr Maria Wells (Executive Officer – Research) by emailing [M.Wells@bath.ac.uk](mailto:M.Wells@bath.ac.uk) or by post to:

Vice-Chancellor's Office, University of Bath, Claverton Down, Bath, BA2 7AY, United Kingdom

**Where else can I access support?**

If after completing this survey you feel distressed and are worried about your emotional well-being, we recommend that you contact your General Practitioner (GP) to discuss this.

***The following list of organisations and publications may be able to provide you with help, both practical and in terms of understanding how you are feeling:***

Child Bereavement UK

Child Bereavement UK is the UK's leading organisation that supports families when a baby or child of any age dies or is dying or when a child is facing bereavement.

Website: [www.childbereavementuk.org](http://www.childbereavementuk.org)  
Helpline: 0800 02 888 40

Sands (Stillbirth and neonatal death charity)

Sands is a national charity that offers support to anyone affected by the death of a baby.

Website: [www.sands.org.uk](http://www.sands.org.uk)  
Helpline: 020 7436 5881

Miscarriage Association

Miscarriage Association is a national organisation that offers support to anyone who has been affected by miscarriage.

Website: [www.miscarriageassociation.org.uk](http://www.miscarriageassociation.org.uk)  
Helpline: 01924 200 799

Birth Trauma Association (BTA)

Birth Trauma Association supports women who have had a traumatic birth experience.

Website: [www.birthtraumaassociation.org.uk](http://www.birthtraumaassociation.org.uk)

Cruse Bereavement Care

Cruse Bereavement Care promotes the wellbeing of bereaved people and supports anyone bereaved by death to understand their grief and cope with loss.

Website: [www.cruse.org.uk](http://www.cruse.org.uk)  
Helpline: 0808 808 1677

Care for the Family

Care for the family is a national charity which aims to promote strong family life and to help those who face family difficulties, including bereavement.

Website: [careforfamily.org.uk](http://careforfamily.org.uk)

Samaritans

Available 24 hours a day to provide confidential emotional support for people who are experiencing feelings of distress or despair.

Website: [samaritans.org.uk](http://samaritans.org.uk)  
Helpline: 116 123



## EMOTIONAL REACTIONS TO LOSS IN PREGNANCY

### *Demographic Questionnaire*

Please answer the following questions about yourself. This is to let us know whether the survey has reached a wide range of different people and to check that you are eligible for the study.

#### About you

##### **1. How would you describe your gender?**

- ☐ Female ☐ Other  
☐ Male

If you selected Other, please specify if you wish (optional)

\_\_\_\_\_

##### **2. How would you describe your ethnicity?** Choose one option that describes your ethnic group best.

- ☐ White British ☐ Black British  
☐ Other White background ☐ Other Black  
☐ Asian British ☐ Multiple/Mixed Ethnic Groups  
☐ Other Asian ☐ Other

If you selected Other, please specify if you wish (optional)

\_\_\_\_\_

##### **3. How would you describe your marital status?**

- ☐ Single, never married ☐ Divorced or separated  
☐ Married/civil partnership or cohabiting ☐ Prefer not to say

##### **4. What is the highest level of schooling you have completed?**

- ☐ No schooling ☐ Degree qualification or above  
☐ Left school without qualifications ☐ Prefer not to say  
☐ GCSE qualifications or equivalent ☐ Other  
☐ A-Level or equivalent

If you selected Other, please specify if you wish (optional)

\_\_\_\_\_

**5. What is your current employment status?**

- |  |                                    |
|--|------------------------------------|
| <input type="checkbox"/> Full-time         | <input type="checkbox"/> Student   |
| <input type="checkbox"/> Part-time         | <input type="checkbox"/> Homemaker |
| <input type="checkbox"/> Prefer not to say | <input type="checkbox"/> Other     |

If you selected Other, please specify if you wish (optional)

\_\_\_\_\_

**6. What is your current profession (if applicable)?**

\_\_\_\_\_  
\_\_\_\_\_

**7. How old are you?**

\_\_\_\_\_  
\_\_\_\_\_

About your pregnancy

**8b) Have you experienced a stillbirth (at 24 weeks of pregnancy or later) or an early miscarriage (before 12 weeks of pregnancy) between 24 months and 6 years ago?**

If you have experienced multiple miscarriages or stillbirths in this time, please answer this question in relation to the loss that affected you most. There is no right or wrong choice for this. We ask you to focus on a particular loss to help answer the questions as they are worded. For any other losses in this time period, please outline them in question 9.

- |  |                             |
|--|-----------------------------|
| <input type="checkbox"/> Yes, a stillbirth         | <input type="checkbox"/> No |
| <input type="checkbox"/> Yes, an early miscarriage |                             |

**8c) If you answered "Yes" in question 8b), what was the exact date of your stillbirth or early miscarriage?**

\_\_\_\_\_

**8d) Approximately how many weeks pregnant were you when you lost this baby?**

- |                                      |  |
|--------------------------------------|--|
| <input type="checkbox"/> 1-4 weeks   | <input type="checkbox"/> 28-31 weeks   |
| <input type="checkbox"/> 5-8 weeks   | <input type="checkbox"/> 32-35 weeks   |
| <input type="checkbox"/> 9-12 weeks  | <input type="checkbox"/> 36-40 weeks   |
| <input type="checkbox"/> 24-27 weeks | <input type="checkbox"/> Over 40 weeks |

**9. Previous losses**

**Have you had other miscarriages or stillbirths in the past? If so, when?** Please answer this question in relation to any other losses you have experienced, other than the one in question 8b) to 8d) above.

**a) Early miscarriage** (occurring in the 0-12 week period of your pregnancy)

- ☐ Yes ☐ No

Dates (year) \_\_\_\_\_

**b) Late miscarriage** (occurring in the 13-23 week period of your pregnancy)

- ☐ Yes ☐ No

Dates (year) \_\_\_\_\_

**c) Stillbirth** (occurring at 24 weeks of pregnancy or later)

- ☐ Yes ☐ No

Dates (year) \_\_\_\_\_

**10. Do you have any living children?**

- ☐ Yes ☐ Prefer not to say
- ☐ No

If you selected Yes, please specify how old your children are and whether they are children that you gave birth to, adopted, fostered or your step-children.

\_\_\_\_\_  
\_\_\_\_\_

**11. Are you pregnant at the moment?**

- ☐ Yes ☐ No ☐ Prefer not to say

Thank you for answering the questions about yourself and your loss. The next questions will ask you about your emotional wellbeing. Please be as honest as you can and remember that there are no right or wrong answers.

## EMOTIONAL REACTIONS TO LOSS IN PREGNANCY

### *Survey Questions*

Miscarriage and stillbirth are unique painful experiences and each person will experience different emotional reactions. As time passes, some people will find their emotional reactions more manageable, but for others they can last longer and be very distressing.

Please answer the following questions about your emotional well-being. The questions will ask about different reactions and some may not be something that you have experienced. The questions are from standardised questionnaires that have been tested in different studies – using them means that the results from this project will be more reliable. However, this does mean that the wording may not always seem appropriate. **For all questions that refer to “the event”, “the trauma” or “the incident”, please think about “losing my baby”. If you have experienced multiple losses in the last 24 months to 6 years, please answer the questions in relation to the loss that affected you most. There is no right or wrong choice for this. We ask you to focus on a particular loss to help answer the questions as they are worded.**

Please do not spend very long thinking about any of your answers – read each question and answer it as quickly as possible. Try to be as honest as you can and remember that there are no right or wrong answers.

The PHQ-8, GAD-7, PGS, PCL-5, PTGI, CBI, DTQ and ERRI questionnaires have not been included in this appendix in order to comply with copyright law. Only questionnaires relating to actual self-disclosure and experience of completing the survey have been included.



### Talking to Others

If you have experienced multiple losses in the last 24 months to 6 years, please answer the following questions in relation to the loss that affected you most.

#### 1. First month

a) In the first month after losing your baby, how much time did you spend talking to others (e.g. friends, family, healthcare staff or anyone else) about your feelings about losing your baby? (Please give an estimation of the number of hours)

---

b) In the first month after losing your baby, do you think you talked enough about your feelings about losing your baby?

No, I did not talk enough

Yes, it was about right

No, I talked too much

☐☐☐

#### 2. First Year

a) In the first year after losing your baby, how much time did you spend talking to others (e.g. friends, family, healthcare staff or anyone else) about your feelings about losing your baby? (Please give an estimation of the number of hours)

---

b) In the first year after losing your baby, do you think you talked enough about your feelings about losing your baby?

No, I did not talk enough

Yes, it was about right

No, I talked too much

☐☐☐

#### 3. Second Year

a) In the second year after losing your baby, how much time did you spend talking to others (e.g. friends, family, healthcare staff or anyone else) about your feelings about losing your baby? (Please give an estimation of the number of hours)

---

b) In the second year after losing your baby, do you think you talked enough about your feelings about losing your baby?

No, I did not talk enough

Yes, it was about right

No, I talked too much

☐☐☐

Thank you for taking part in this study. We really value your participation. We hope that the outcomes of this project will improve our understanding of emotional reactions following early miscarriage and stillbirth and improve the help and support offered to women and their families in the future.

#### QUESTIONS ABOUT COMPLETING THE SURVEY

**Has your experience of completing these questionnaires at the moment been:**

- |   |  |
|---|--|
| <input type="checkbox"/> Very positive                | <input type="checkbox"/> Negative          |
| <input type="checkbox"/> Positive                     | <input type="checkbox"/> Very negative     |
| <input type="checkbox"/> Neither positive or negative | <input type="checkbox"/> Prefer not to say |

**Do you believe that completing these questionnaires will be a positive or negative experience for you in the long-term?**

- |   |  |
|---|--|
| <input type="checkbox"/> Very positive                | <input type="checkbox"/> Negative          |
| <input type="checkbox"/> Positive                     | <input type="checkbox"/> Very negative     |
| <input type="checkbox"/> Neither positive or negative | <input type="checkbox"/> Prefer not to say |

**Do you think that you may contact someone (either your GP or another professional organisation/support group) to speak about your loss as a result of taking part in this research?**

- |                              |  |
|------------------------------|--|
| <input type="checkbox"/> Yes | <input type="checkbox"/> Not sure          |
| <input type="checkbox"/> No  | <input type="checkbox"/> Prefer not to say |

***Having completed this survey, I agree to submit my responses and understand that once I have submitted the survey I will be unable to ask for my data to be withdrawn as it will not be possible to identify my anonymous responses. I agree to the research team keeping and processing the data I have provided during the course of this study and archiving and sharing this data with other researchers in the future.***

- |                              |                             |
|------------------------------|-----------------------------|
| <input type="checkbox"/> Yes | <input type="checkbox"/> No |
|------------------------------|-----------------------------|

## EMOTIONAL REACTIONS TO LOSS IN PREGNANCY

### *Debrief Sheet*

Thank you for taking part in this study. We really value your participation and hope that the outcomes of this project will improve our understanding and enable us to improve the help and support offered to women and their families following early miscarriage and stillbirth.

#### **What happens next?**

On completion of the project, a summary of the findings will be made available to participants who wish to receive them in the summer of 2018. If you would like to receive this written summary of the findings, please email the research team at [erlp@bath.ac.uk](mailto:erlp@bath.ac.uk) to request this.

#### **What do I do if I have concerns about the project?**

Every care has been taken to ensure your safety during the course of the study. If you have any concerns about the project or your participation in the research, please contact a member of the research team by emailing [erlp@bath.ac.uk](mailto:erlp@bath.ac.uk). If you wish, we can arrange for someone to call you to discuss your concerns within three working days.

#### **Where can I access support?**

We recommend that you contact your General Practitioner (GP) if you feel distressed and are worried about your emotional well-being after taking part in this study. If you have found it upsetting to answer questions about your experiences, you may wish to seek some support.

We have prepared a list of resources for participants who have taken part in the project so that you are aware of what options are available if you feel you would like some additional support. Please note that we do not expect you to be looking for support but we wanted to keep this list as comprehensive as possible. The list is not exhaustive however and we cannot specifically recommend each website/organisation or guarantee the quality of each.

The following list of organisations may be able to provide you with help, both practical and in terms of understanding how you are feeling:

#### **Child Bereavement UK**

Child Bereavement UK is the UK's leading organisation that supports families when a baby or child of any age dies or is dying or when a child is facing bereavement.

Website: [www.childbereavementuk.org](http://www.childbereavementuk.org)  
Helpline: 0800 02 888 40

#### **Sands (Stillbirth and neonatal death charity)**

Sands is a national charity that offers support to anyone affected by the death of a baby.

Website: [www.sands.org.uk](http://www.sands.org.uk)  
Helpline: 020 7436 5881

**Miscarriage Association**

Miscarriage Association is a national organisation that offers support to anyone who has been affected by miscarriage.

*Website:* [www.miscarriageassociation.org.uk](http://www.miscarriageassociation.org.uk)  
*Helpline:* 01924 200 799

**Birth Trauma Association (BTA)**

Birth Trauma Association supports women who have had a traumatic birth experience.

*Website:* [www.birthtraumaassociation.org.uk](http://www.birthtraumaassociation.org.uk)

**Cruse Bereavement Care**

Cruse Bereavement Care promotes the wellbeing of bereaved people and supports anyone bereaved by death to understand their grief and cope with loss.

*Website:* [www.cruse.org.uk](http://www.cruse.org.uk)  
*Helpline:* 0808 808 1677

**Care for the Family**

Care for the family is a national charity which aims to promote strong family life and to help those who face family difficulties, including bereavement.

*Website:* [careforfamily.org.uk](http://careforfamily.org.uk)

**Samaritans**

Available 24 hours a day to provide confidential emotional support for people who are experiencing feelings of distress or despair.

*Website:* [samaritans.org.uk](http://samaritans.org.uk)  
*Helpline:* 116 123